






# 4 Integration



The area of a parabolic region can be approximated as the sum of the areas of rectangles. As you increase the number of rectangles, the approximation tends to become more and more accurate. In Section 4.2, you will learn how the limit process can be used to find areas of a wide variety of regions. This process is called *integration* and is closely related to differentiation.

This photo of a jet breaking the sound barrier was taken by Ensign John Gay. At the time the photo was taken, was the jet's velocity constant or changing? Why?



©Corbis Sygma

## Section 4.1

## Antiderivatives and Indefinite Integration

- Write the general solution of a differential equation.
- Use indefinite integral notation for antiderivatives.
- Use basic integration rules to find antiderivatives.
- Find a particular solution of a differential equation.

## EXPLORATION

**Finding Antiderivatives** For each derivative, describe the original function  $F$ .

a.  $F'(x) = 2x$

b.  $F'(x) = x$

c.  $F'(x) = x^2$

d.  $F'(x) = \frac{1}{x^2}$

e.  $F'(x) = \frac{1}{x^3}$

f.  $F'(x) = \cos x$

What strategy did you use to find  $F$ ?

## Antiderivatives

Suppose you were asked to find a function  $F$  whose derivative is  $f(x) = 3x^2$ . From your knowledge of derivatives, you would probably say that

$$F(x) = x^3 \text{ because } \frac{d}{dx}[x^3] = 3x^2.$$

The function  $F$  is an *antiderivative* of  $f$ .

## Definition of an Antiderivative

A function  $F$  is an **antiderivative** of  $f$  on an interval  $I$  if  $F'(x) = f(x)$  for all  $x$  in  $I$ .

Note that  $F$  is called *an* antiderivative of  $f$ , rather than *the* antiderivative of  $f$ . To see why, observe that

$$F_1(x) = x^3, \quad F_2(x) = x^3 - 5, \quad \text{and} \quad F_3(x) = x^3 + 97$$

are all antiderivatives of  $f(x) = 3x^2$ . In fact, for any constant  $C$ , the function given by  $F(x) = x^3 + C$  is an antiderivative of  $f$ .

## THEOREM 4.1 Representation of Antiderivatives

If  $F$  is an antiderivative of  $f$  on an interval  $I$ , then  $G$  is an antiderivative of  $f$  on the interval  $I$  if and only if  $G$  is of the form  $G(x) = F(x) + C$ , for all  $x$  in  $I$  where  $C$  is a constant.

**Proof** The proof of Theorem 4.1 in one direction is straightforward. That is, if  $G(x) = F(x) + C$ ,  $F'(x) = f(x)$ , and  $C$  is a constant, then

$$G'(x) = \frac{d}{dx}[F(x) + C] = F'(x) + 0 = f(x).$$

To prove this theorem in the other direction, assume that  $G$  is an antiderivative of  $f$ . Define a function  $H$  such that

$$H(x) = G(x) - F(x).$$

If  $H$  is not constant on the interval  $I$ , there must exist  $a$  and  $b$  ( $a < b$ ) in the interval such that  $H(a) \neq H(b)$ . Moreover, because  $H$  is differentiable on  $(a, b)$ , you can apply the Mean Value Theorem to conclude that there exists some  $c$  in  $(a, b)$  such that

$$H'(c) = \frac{H(b) - H(a)}{b - a}.$$

Because  $H(b) \neq H(a)$ , it follows that  $H'(c) \neq 0$ . However, because  $G'(c) = F'(c)$ , you know that  $H'(c) = G'(c) - F'(c) = 0$ , which contradicts the fact that  $H'(c) \neq 0$ . Consequently, you can conclude that  $H(x)$  is a constant,  $C$ . So,  $G(x) - F(x) = C$  and it follows that  $G(x) = F(x) + C$ .

Using Theorem 4.1, you can represent the entire family of antiderivatives of a function by adding a constant to a *known* antiderivative. For example, knowing that  $D_x[x^2] = 2x$ , you can represent the family of *all* antiderivatives of  $f(x) = 2x$  by

$$G(x) = x^2 + C \quad \text{Family of all antiderivatives of } f(x) = 2x$$

where  $C$  is a constant. The constant  $C$  is called the **constant of integration**. The family of functions represented by  $G$  is the **general antiderivative** of  $f$ , and  $G(x) = x^2 + C$  is the **general solution** of the differential equation

$$G'(x) = 2x. \quad \text{Differential equation}$$

A **differential equation** in  $x$  and  $y$  is an equation that involves  $x$ ,  $y$ , and derivatives of  $y$ . For instance,  $y' = 3x$  and  $y' = x^2 + 1$  are examples of differential equations.

### EXAMPLE 1 Solving a Differential Equation

Find the general solution of the differential equation  $y' = 2$ .


**Solution** To begin, you need to find a function whose derivative is 2. One such function is

$$y = 2x. \quad 2x \text{ is an antiderivative of } 2.$$

Now, you can use Theorem 4.1 to conclude that the general solution of the differential equation is

$$y = 2x + C. \quad \text{General solution}$$

The graphs of several functions of the form  $y = 2x + C$  are shown in Figure 4.1.



Functions of the form  $y = 2x + C$   
Figure 4.1

### Notation for Antiderivatives

When solving a differential equation of the form

$$\frac{dy}{dx} = f(x)$$

it is convenient to write it in the equivalent differential form

$$dy = f(x) dx.$$

The operation of finding all solutions of this equation is called **antidifferentiation** (or **indefinite integration**) and is denoted by an integral sign  $\int$ . The general solution is denoted by

$$y = \int f(x) dx = F(x) + C.$$

Variable of integration     Constant of integration  
Integrand

**NOTE** In this text, the notation  $\int f(x) dx = F(x) + C$  means that  $F$  is an antiderivative of  $f$  on an interval.

The expression  $\int f(x) dx$  is read as the *antiderivative of  $f$  with respect to  $x$* . So, the differential  $dx$  serves to identify  $x$  as the variable of integration. The term **indefinite integral** is a synonym for antiderivative.

## Basic Integration Rules

The inverse nature of integration and differentiation can be verified by substituting  $F'(x)$  for  $f(x)$  in the indefinite integration definition to obtain

$$\int F'(x) dx = F(x) + C.$$

Integration is the “inverse” of differentiation.

Moreover, if  $\int f(x) dx = F(x) + C$ , then

$$\frac{d}{dx} \left[ \int f(x) dx \right] = f(x).$$

Differentiation is the “inverse” of integration.

These two equations allow you to obtain integration formulas directly from differentiation formulas, as shown in the following summary.

### Basic Integration Rules

#### Differentiation Formula

$$\frac{d}{dx}[C] = 0$$

$$\frac{d}{dx}[kx] = k$$

$$\frac{d}{dx}[kf(x)] = kf'(x)$$

$$\frac{d}{dx}[f(x) \pm g(x)] = f'(x) \pm g'(x)$$

$$\frac{d}{dx}[x^n] = nx^{n-1}$$

$$\frac{d}{dx}[\sin x] = \cos x$$

$$\frac{d}{dx}[\cos x] = -\sin x$$

$$\frac{d}{dx}[\tan x] = \sec^2 x$$

$$\frac{d}{dx}[\sec x] = \sec x \tan x$$

$$\frac{d}{dx}[\cot x] = -\csc^2 x$$

$$\frac{d}{dx}[\csc x] = -\csc x \cot x$$

#### Integration Formula

$$\int 0 dx = C$$

$$\int k dx = kx + C$$

$$\int kf(x) dx = k \int f(x) dx$$

$$\int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1 \quad \text{Power Rule}$$

$$\int \cos x dx = \sin x + C$$

$$\int \sin x dx = -\cos x + C$$

$$\int \sec^2 x dx = \tan x + C$$

$$\int \sec x \tan x dx = \sec x + C$$

$$\int \csc^2 x dx = -\cot x + C$$

$$\int \csc x \cot x dx = -\csc x + C$$

**NOTE** Note that the Power Rule for Integration has the restriction that  $n \neq -1$ . The evaluation of  $\int 1/x dx$  must wait until the introduction of the natural logarithm function in Chapter 5.

**EXAMPLE 2** Applying the Basic Integration RulesDescribe the antiderivatives of  $3x$ .

**Solution**

$$\begin{aligned}
 \int 3x \, dx &= 3 \int x \, dx && \text{Constant Multiple Rule} \\
 &= 3 \int x^1 \, dx && \text{Rewrite } x \text{ as } x^1. \\
 &= 3 \left( \frac{x^2}{2} \right) + C && \text{Power Rule } (n = 1) \\
 &= \frac{3}{2} x^2 + C && \text{Simplify.}
 \end{aligned}$$

So, the antiderivatives of  $3x$  are of the form  $\frac{3}{2}x^2 + C$ , where  $C$  is any constant.

When indefinite integrals are evaluated, a strict application of the basic integration rules tends to produce complicated constants of integration. For instance, in Example 2, you could have written

$$\int 3x \, dx = 3 \int x \, dx = 3 \left( \frac{x^2}{2} + C \right) = \frac{3}{2} x^2 + 3C.$$

However, because  $C$  represents *any* constant, it is both cumbersome and unnecessary to write  $3C$  as the constant of integration. So,  $\frac{3}{2}x^2 + 3C$  is written in the simpler form,  $\frac{3}{2}x^2 + C$ .

In Example 2, note that the general pattern of integration is similar to that of differentiation.

Original integral



Rewrite



Integrate



Simplify

**EXAMPLE 3** Rewriting Before Integrating

**TECHNOLOGY** Some software programs, such as *Derive*, *Maple*, *Mathcad*, *Mathematica*, and the *TI-89*, are capable of performing integration symbolically. If you have access to such a symbolic integration utility, try using it to evaluate the indefinite integrals in Example 3.

Original Integral	Rewrite	Integrate	Simplify
a. $\int \frac{1}{x^3} \, dx$	$\int x^{-3} \, dx$	$\frac{x^{-2}}{-2} + C$	$-\frac{1}{2x^2} + C$
b. $\int \sqrt{x} \, dx$	$\int x^{1/2} \, dx$	$\frac{x^{3/2}}{3/2} + C$	$\frac{2}{3} x^{3/2} + C$
c. $\int 2 \sin x \, dx$	$2 \int \sin x \, dx$	$2(-\cos x) + C$	$-2 \cos x + C$

Remember that you can check your answer to an antidifferentiation problem by differentiating. For instance, in Example 3(b), you can check that  $\frac{2}{3}x^{3/2} + C$  is the correct antiderivative by differentiating the answer to obtain

$$D_x \left[ \frac{2}{3} x^{3/2} + C \right] = \left( \frac{2}{3} \right) \left( \frac{3}{2} \right) x^{1/2} = \sqrt{x}. \quad \text{Use differentiation to check antiderivative.}$$



indicates that in the HM mathSpace® CD-ROM and the online Eduspace® system for this text, you will find an Open Exploration, which further explores this example using the computer algebra systems Maple, Mathcad, Mathematica, and Derive.

The basic integration rules listed earlier in this section allow you to integrate any polynomial function, as shown in Example 4.

#### EXAMPLE 4 Integrating Polynomial Functions

$$\begin{aligned}\text{a. } \int dx &= \int 1 \, dx \\ &= x + C\end{aligned}$$

Integrand is understood to be 1.

Integrate.

$$\begin{aligned}\text{b. } \int (x + 2) \, dx &= \int x \, dx + \int 2 \, dx \\ &= \frac{x^2}{2} + C_1 + 2x + C_2 \\ &= \frac{x^2}{2} + 2x + C\end{aligned}$$

Integrate.

$C = C_1 + C_2$

The second line in the solution is usually omitted.

$$\begin{aligned}\text{c. } \int (3x^4 - 5x^2 + x) \, dx &= 3\left(\frac{x^5}{5}\right) - 5\left(\frac{x^3}{3}\right) + \frac{x^2}{2} + C \\ &= \frac{3}{5}x^5 - \frac{5}{3}x^3 + \frac{1}{2}x^2 + C\end{aligned}$$

Integrate.

Simplify.

#### EXAMPLE 5 Rewriting Before Integrating

$$\begin{aligned}\int \frac{x+1}{\sqrt{x}} \, dx &= \int \left( \frac{x}{\sqrt{x}} + \frac{1}{\sqrt{x}} \right) \, dx \\ &= \int (x^{1/2} + x^{-1/2}) \, dx \\ &= \frac{x^{3/2}}{3/2} + \frac{x^{1/2}}{1/2} + C \\ &= \frac{2}{3}x^{3/2} + 2x^{1/2} + C \\ &= \frac{2}{3}\sqrt{x}(x+3) + C\end{aligned}$$

Rewrite as two fractions.

Rewrite with fractional exponents.

Integrate.

Simplify.

**NOTE** When integrating quotients, do not integrate the numerator and denominator separately. This is no more valid in integration than it is in differentiation. For instance, in Example 5, be sure you understand that

$$\int \frac{x+1}{\sqrt{x}} \, dx = \frac{2}{3}\sqrt{x}(x+3) + C \text{ is not the same as } \frac{\int (x+1) \, dx}{\int \sqrt{x} \, dx} = \frac{\frac{1}{2}x^2 + x + C_1}{\frac{2}{3}x\sqrt{x} + C_2}.$$

#### EXAMPLE 6 Rewriting Before Integrating


$$\begin{aligned}\int \frac{\sin x}{\cos^2 x} \, dx &= \int \left( \frac{1}{\cos x} \right) \left( \frac{\sin x}{\cos x} \right) \, dx \\ &= \int \sec x \tan x \, dx \\ &= \sec x + C\end{aligned}$$

Rewrite as a product.

Rewrite using trigonometric identities.

Integrate.





The particular solution that satisfies the initial condition  $F(2) = 4$  is  $F(x) = x^3 - x - 2$ .

Figure 4.2

## Initial Conditions and Particular Solutions

You have already seen that the equation  $y = \int f(x) dx$  has many solutions (each differing from the others by a constant). This means that the graphs of any two antiderivatives of  $f$  are vertical translations of each other. For example, Figure 4.2 shows the graphs of several antiderivatives of the form

$$y = \int (3x^2 - 1) dx = x^3 - x + C \quad \text{General solution}$$

for various integer values of  $C$ . Each of these antiderivatives is a solution of the differential equation

$$\frac{dy}{dx} = 3x^2 - 1.$$

In many applications of integration, you are given enough information to determine a **particular solution**. To do this, you need only know the value of  $y = F(x)$  for one value of  $x$ . This information is called an **initial condition**. For example, in Figure 4.2, only one curve passes through the point  $(2, 4)$ . To find this curve, you can use the following information.

$$F(x) = x^3 - x + C \quad \text{General solution}$$

$$F(2) = 4 \quad \text{Initial condition}$$

By using the initial condition in the general solution, you can determine that  $F(2) = 8 - 2 + C = 4$ , which implies that  $C = -2$ . So, you obtain

$$F(x) = x^3 - x - 2. \quad \text{Particular solution}$$

## EXAMPLE 7 Finding a Particular Solution

Find the general solution of

$$F'(x) = \frac{1}{x^2}, \quad x > 0$$

and find the particular solution that satisfies the initial condition  $F(1) = 0$ .

**Solution** To find the general solution, integrate to obtain

$$F(x) = \int \frac{1}{x^2} dx \quad F(x) = \int F'(x) dx$$

$$= \int x^{-2} dx \quad \text{Rewrite as a power.}$$

$$= \frac{x^{-1}}{-1} + C \quad \text{Integrate.}$$


$$= -\frac{1}{x} + C, \quad x > 0. \quad \text{General solution}$$

Using the initial condition  $F(1) = 0$ , you can solve for  $C$  as follows.

$$F(1) = -\frac{1}{1} + C = 0 \quad \Rightarrow \quad C = 1$$

So, the particular solution, as shown in Figure 4.3, is

$$F(x) = -\frac{1}{x} + 1, \quad x > 0. \quad \text{Particular solution}$$



The particular solution that satisfies the initial condition  $F(1) = 0$  is  $F(x) = -(1/x) + 1, x > 0$ .

Figure 4.3

So far in this section you have been using  $x$  as the variable of integration. In applications, it is often convenient to use a different variable. For instance, in the following example involving *time*, the variable of integration is  $t$ .

### EXAMPLE 8 Solving a Vertical Motion Problem

A ball is thrown upward with an initial velocity of 64 feet per second from an initial height of 80 feet.

- Find the position function giving the height  $s$  as a function of the time  $t$ .
- When does the ball hit the ground?

#### Solution

- Let  $t = 0$  represent the initial time. The two given initial conditions can be written as follows.

$$s(0) = 80$$

Initial height is 80 feet.

$$s'(0) = 64$$

Initial velocity is 64 feet per second.

Using  $-32$  feet per second per second as the acceleration due to gravity, you can write

$$s''(t) = -32$$

$$s'(t) = \int s''(t) dt = \int -32 dt = -32t + C_1.$$

Using the initial velocity, you obtain  $s'(0) = 64 = -32(0) + C_1$ , which implies that  $C_1 = 64$ . Next, by integrating  $s'(t)$ , you obtain

$$s(t) = \int s'(t) dt = \int (-32t + 64) dt = -16t^2 + 64t + C_2.$$

Using the initial height, you obtain

$$s(0) = 80 = -16(0^2) + 64(0) + C_2$$


which implies that  $C_2 = 80$ . So, the position function is

$$s(t) = -16t^2 + 64t + 80. \quad \text{See Figure 4.4.}$$

- Using the position function found in part (a), you can find the time that the ball hits the ground by solving the equation  $s(t) = 0$ .

$$\begin{aligned} s(t) &= -16t^2 + 64t + 80 = 0 \\ -16(t + 1)(t - 5) &= 0 \\ t &= -1, 5 \end{aligned}$$

Because  $t$  must be positive, you can conclude that the ball hits the ground 5 seconds after it was thrown.



Height of a ball at time  $t$

Figure 4.4

**NOTE** In Example 8, note that the position function has the form

$$s(t) = \frac{1}{2}gt^2 + v_0t + s_0$$

where  $g = -32$ ,  $v_0$  is the initial velocity, and  $s_0$  is the initial height, as presented in Section 2.2.

Example 8 shows how to use calculus to analyze vertical motion problems in which the acceleration is determined by a gravitational force. You can use a similar strategy to analyze other linear motion problems (vertical or horizontal) in which the acceleration (or deceleration) is the result of some other force, as you will see in Exercises 77–86.



Before you begin the exercise set, be sure you realize that one of the most important steps in integration is *rewriting the integrand* in a form that fits the basic integration rules. To illustrate this point further, here are some additional examples.

<u>Original Integral</u>	<u>Rewrite</u>	<u>Integrate</u>	<u>Simplify</u>
$\int \frac{2}{\sqrt{x}} dx$	$2 \int x^{-1/2} dx$	$2 \left( \frac{x^{1/2}}{1/2} \right) + C$	$4x^{1/2} + C$
$\int (t^2 + 1)^2 dt$	$\int (t^4 + 2t^2 + 1) dt$	$\frac{t^5}{5} + 2 \left( \frac{t^3}{3} \right) + t + C$	$\frac{1}{5}t^5 + \frac{2}{3}t^3 + t + C$
$\int \frac{x^3 + 3}{x^2} dx$	$\int (x + 3x^{-2}) dx$	$\frac{x^2}{2} + 3 \left( \frac{x^{-1}}{-1} \right) + C$	$\frac{1}{2}x^2 - \frac{3}{x} + C$
$\int \sqrt[3]{x}(x - 4) dx$	$\int (x^{4/3} - 4x^{1/3}) dx$	$\frac{x^{7/3}}{7/3} - 4 \left( \frac{x^{4/3}}{4/3} \right) + C$	$\frac{3}{7}x^{7/3} - 3x^{4/3}$

### Exercises for Section 4.1

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.

In Exercises 1–4, verify the statement by showing that the derivative of the right side equals the integrand of the left side.

- $\int \left( -\frac{9}{x^4} \right) dx = \frac{3}{x^3} + C$
- $\int \left( 4x^3 - \frac{1}{x^2} \right) dx = x^4 + \frac{1}{x} + C$
- $\int (x - 2)(x + 2) dx = \frac{1}{3}x^3 - 4x + C$
- $\int \frac{x^2 - 1}{x^{3/2}} dx = \frac{2(x^2 + 3)}{3\sqrt{x}} + C$

In Exercises 5–8, find the general solution of the differential equation and check the result by differentiation.

- $\frac{dy}{dt} = 3t^2$
- $\frac{dr}{d\theta} = \pi$
- $\frac{dy}{dx} = x^{3/2}$
- $\frac{dy}{dx} = 2x^{-3}$

In Exercises 9–14, complete the table.

<u>Original Integral</u>	<u>Rewrite</u>	<u>Integrate</u>	<u>Simplify</u>
9. $\int \sqrt[3]{x} dx$			
10. $\int \frac{1}{x^2} dx$			
11. $\int \frac{1}{x\sqrt{x}} dx$			
12. $\int x(x^2 + 3) dx$			
13. $\int \frac{1}{2x^3} dx$			
14. $\int \frac{1}{(3x)^2} dx$			





In Exercises 15–34, find the indefinite integral and check the result by differentiation.

- $\int (x + 3) dx$
- $\int (5 - x) dx$
- $\int (2x - 3x^2) dx$
- $\int (4x^3 + 6x^2 - 1) dx$
- $\int (x^3 + 2) dx$
- $\int (x^3 - 4x + 2) dx$
- $\int (x^{3/2} + 2x + 1) dx$
- $\int \left( \sqrt{x} + \frac{1}{2\sqrt{x}} \right) dx$
- $\int \sqrt[3]{x^2} dx$
- $\int (\sqrt[4]{x^3} + 1) dx$
- $\int \frac{1}{x^3} dx$
- $\int \frac{1}{x^4} dx$
- $\int \frac{x^2 + x + 1}{\sqrt{x}} dx$
- $\int \frac{x^2 + 2x - 3}{x^4} dx$
- $\int (x + 1)(3x - 2) dx$
- $\int (2t^2 - 1)^2 dt$
- $\int y^2 \sqrt{y} dy$
- $\int (1 + 3t)t^2 dt$
- $\int dx$
- $\int 3 dt$

In Exercises 35–42, find the indefinite integral and check the result by differentiation.


- $\int (2 \sin x + 3 \cos x) dx$
- $\int (t^2 - \sin t) dt$
- $\int (1 - \csc t \cot t) dt$
- $\int (\theta^2 + \sec^2 \theta) d\theta$
- $\int (\sec^2 \theta - \sin \theta) d\theta$
- $\int \sec y (\tan y - \sec y) dy$
- $\int (\tan^2 y + 1) dy$
- $\int \frac{\cos x}{1 - \cos^2 x} dx$

In Exercises 43–46, the graph of the derivative of a function is given. Sketch the graphs of *two* functions that have the given derivative. (There is more than one correct answer.) To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).





In Exercises 47 and 48, find the equation for  $y$ , given the derivative and the indicated point on the curve.

47.  $\frac{dy}{dx} = 2x - 1$




48.  $\frac{dy}{dx} = 2(x - 1)$




 **Slope Fields** In Exercises 49–52, a differential equation, a point, and a slope field are given. A *slope field* (or *direction field*) consists of line segments with slopes given by the differential equation. These line segments give a visual perspective of the slopes of the solutions of the differential equation. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the indicated point. (To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).) (b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution. Compare the result with the sketches in part (a).


49.  $\frac{dy}{dx} = \frac{1}{2}x - 1, (4, 2)$




50.  $\frac{dy}{dx} = x^2 - 1, (-1, 3)$




51.  $\frac{dy}{dx} = \cos x, (0, 4)$



52.  $\frac{dy}{dx} = -\frac{1}{x^2}, x > 0, (1, 3)$



 **Slope Fields** In Exercises 53 and 54, (a) use a graphing utility to graph a slope field for the differential equation, (b) use integration and the given point to find the particular solution of the differential equation, and (c) graph the solution and the slope field in the same viewing window.

53.  $\frac{dy}{dx} = 2x, (-2, -2)$

54.  $\frac{dy}{dx} = 2\sqrt{x}, (4, 12)$

In Exercises 55–62, solve the differential equation.

55.  $f'(x) = 4x, f(0) = 6$

56.  $g'(x) = 6x^2, g(0) = -1$

57.  $h'(t) = 8t^3 + 5, h(1) = -4$

58.  $f'(s) = 6s - 8s^3, f(2) = 3$

59.  $f''(x) = 2, f'(2) = 5, f(2) = 10$

60.  $f''(x) = x^2, f'(0) = 6, f(0) = 3$

61.  $f''(x) = x^{-3/2}, f'(4) = 2, f(0) = 0$

62.  $f''(x) = \sin x, f'(0) = 1, f(0) = 6$

63. **Tree Growth** An evergreen nursery usually sells a certain shrub after 6 years of growth and shaping. The growth rate during those 6 years is approximated by  $dh/dt = 1.5t + 5$ , where  $t$  is the time in years and  $h$  is the height in centimeters. The seedlings are 12 centimeters tall when planted ( $t = 0$ ).

(a) Find the height after  $t$  years.

(b) How tall are the shrubs when they are sold?

64. **Population Growth** The rate of growth  $dP/dt$  of a population of bacteria is proportional to the square root of  $t$ , where  $P$  is the population size and  $t$  is the time in days ( $0 \leq t \leq 10$ ). That is,  $dP/dt = k\sqrt{t}$ . The initial size of the population is 500. After 1 day the population has grown to 600. Estimate the population after 7 days.

### Writing About Concepts

65. Use the graph of  $f'$  shown in the figure to answer the following, given that  $f(0) = -4$ .
- Approximate the slope of  $f$  at  $x = 4$ . Explain.
  - Is it possible that  $f(2) = -1$ ? Explain.
  - Is  $f(5) - f(4) > 0$ ? Explain.
  - Approximate the value of  $x$  where  $f$  is maximum. Explain.
  - Approximate any intervals in which the graph of  $f$  is concave upward and any intervals in which it is concave downward. Approximate the  $x$ -coordinates of any points of inflection.
  - Approximate the  $x$ -coordinate of the minimum of  $f''(x)$ .
  - Sketch an approximate graph of  $f$ . To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).




Figure for 65




Figure for 66

66. The graphs of  $f$  and  $f'$  each pass through the origin. Use the graph of  $f''$  shown in the figure to sketch the graphs of  $f$  and  $f'$ . To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).

**Vertical Motion** In Exercises 67–70, use  $a(t) = -32$  feet per second per second as the acceleration due to gravity. (Neglect air resistance.)

- A ball is thrown vertically upward from a height of 6 feet with an initial velocity of 60 feet per second. How high will the ball go?
- Show that the height above the ground of an object thrown upward from a point  $s_0$  feet above the ground with an initial velocity of  $v_0$  feet per second is given by the function  $f(t) = -16t^2 + v_0t + s_0$ .
- With what initial velocity must an object be thrown upward (from ground level) to reach the top of the Washington Monument (approximately 550 feet)?
- A balloon, rising vertically with a velocity of 16 feet per second, releases a sandbag at the instant it is 64 feet above the ground.
  - How many seconds after its release will the bag strike the ground?
  - At what velocity will it hit the ground?

**Vertical Motion** In Exercises 71–74, use  $a(t) = -9.8$  meters per second per second as the acceleration due to gravity. (Neglect air resistance.)

- Show that the height above the ground of an object thrown upward from a point  $s_0$  meters above the ground with an initial velocity of  $v_0$  meters per second is given by the function

$$f(t) = -4.9t^2 + v_0t + s_0.$$

- The Grand Canyon is 1800 meters deep at its deepest point. A rock is dropped from the rim above this point. Write the height of the rock as a function of the time  $t$  in seconds. How long will it take the rock to hit the canyon floor?
- A baseball is thrown upward from a height of 2 meters with an initial velocity of 10 meters per second. Determine its maximum height.
- With what initial velocity must an object be thrown upward (from a height of 2 meters) to reach a maximum height of 200 meters?

- Lunar Gravity** On the moon, the acceleration due to gravity is  $-1.6$  meters per second per second. A stone is dropped from a cliff on the moon and hits the surface of the moon 20 seconds later. How far did it fall? What was its velocity at impact?

- Escape Velocity** The minimum velocity required for an object to escape Earth's gravitational pull is obtained from the solution of the equation

$$\int v \, dv = -GM \int \frac{1}{y^2} \, dy$$

where  $v$  is the velocity of the object projected from Earth,  $y$  is the distance from the center of Earth,  $G$  is the gravitational constant, and  $M$  is the mass of Earth. Show that  $v$  and  $y$  are related by the equation

$$v^2 = v_0^2 + 2GM \left( \frac{1}{y} - \frac{1}{R} \right)$$

where  $v_0$  is the initial velocity of the object and  $R$  is the radius of Earth.

**Rectilinear Motion** In Exercises 77–80, consider a particle moving along the  $x$ -axis where  $x(t)$  is the position of the particle at time  $t$ ,  $x'(t)$  is its velocity, and  $x''(t)$  is its acceleration.

- $x(t) = t^3 - 6t^2 + 9t - 2$ ,  $0 \leq t \leq 5$ 
  - Find the velocity and acceleration of the particle.
  - Find the open  $t$ -intervals on which the particle is moving to the right.
  - Find the velocity of the particle when the acceleration is 0.
- Repeat Exercise 77 for the position function  $x(t) = (t - 1)(t - 3)^2$ ,  $0 \leq t \leq 5$ .
- A particle moves along the  $x$ -axis at a velocity of  $v(t) = 1/\sqrt{t}$ ,  $t > 0$ . At time  $t = 1$ , its position is  $x = 4$ . Find the acceleration and position functions for the particle.

## THE SUM OF THE FIRST 100 INTEGERS

Carl Friedrich Gauss's (1777–1855) teacher asked him to add all the integers from 1 to 100. When Gauss returned with the correct answer after only a few moments, the teacher could only look at him in astounded silence. This is what Gauss did:

$$\begin{array}{r} 1 + 2 + 3 + \cdots + 100 \\ 100 + 99 + 98 + \cdots + 1 \\ \hline 101 + 101 + 101 + \cdots + 101 \\ \hline \frac{100 \times 101}{2} = 5050 \end{array}$$

This is generalized by Theorem 4.2, where

$$\sum_{i=1}^{100} i = \frac{100(101)}{2} = 5050.$$

The following properties of summation can be derived using the associative and commutative properties of addition and the distributive property of addition over multiplication. (In the first property,  $k$  is a constant.)

1.  $\sum_{i=1}^n ka_i = k \sum_{i=1}^n a_i$
2.  $\sum_{i=1}^n (a_i \pm b_i) = \sum_{i=1}^n a_i \pm \sum_{i=1}^n b_i$

The next theorem lists some useful formulas for sums of powers. A proof of this theorem is given in Appendix A.

**THEOREM 4.2** Summation Formulas

1.  $\sum_{i=1}^n c = cn$
2.  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$
3.  $\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$
4.  $\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$

**EXAMPLE 2** Evaluating a Sum

Evaluate  $\sum_{i=1}^n \frac{i+1}{n^2}$  for  $n = 10, 100, 1000$ , and  $10,000$ .

**Solution** Applying Theorem 4.2, you can write

$$\begin{aligned} \sum_{i=1}^n \frac{i+1}{n^2} &= \frac{1}{n^2} \sum_{i=1}^n (i+1) && \text{Factor constant } 1/n^2 \text{ out of sum.} \\ &= \frac{1}{n^2} \left( \sum_{i=1}^n i + \sum_{i=1}^n 1 \right) && \text{Write as two sums.} \\ &= \frac{1}{n^2} \left[ \frac{n(n+1)}{2} + n \right] && \text{Apply Theorem 4.2.} \\ &= \frac{1}{n^2} \left[ \frac{n^2 + 3n}{2} \right] && \text{Simplify.} \\ &= \frac{n+3}{2n} && \text{Simplify.} \end{aligned}$$


Now you can evaluate the sum by substituting the appropriate values of  $n$ , as shown in the table at the left.

$n$	$\sum_{i=1}^n \frac{i+1}{n^2} = \frac{n+3}{2n}$
10	0.65000
100	0.51500
1,000	0.50150
10,000	0.50015

In the table, note that the sum appears to approach a limit as  $n$  increases. Although the discussion of limits at infinity in Section 3.5 applies to a variable  $x$ , where  $x$  can be any real number, many of the same results hold true for limits involving the variable  $n$ , where  $n$  is restricted to positive integer values. So, to find the limit of  $(n+3)/2n$  as  $n$  approaches infinity, you can write


$$\lim_{n \rightarrow \infty} \frac{n+3}{2n} = \frac{1}{2}.$$





Rectangle:  $A = bh$

Figure 4.5




Triangle:  $A = \frac{1}{2}bh$

Figure 4.6


## Area

In Euclidean geometry, the simplest type of plane region is a rectangle. Although people often say that the *formula* for the area of a rectangle is  $A = bh$ , as shown in Figure 4.5, it is actually more proper to say that this is the *definition* of the **area of a rectangle**.


From this definition, you can develop formulas for the areas of many other plane regions. For example, to determine the area of a triangle, you can form a rectangle whose area is twice that of the triangle, as shown in Figure 4.6. Once you know how to find the area of a triangle, you can determine the area of any polygon by subdividing the polygon into triangular regions, as shown in Figure 4.7.



Parallelogram



Hexagon





Polygon

Figure 4.7

Finding the areas of regions other than polygons is more difficult. The ancient Greeks were able to determine formulas for the areas of some general regions (principally those bounded by conics) by the *exhaustion* method. The clearest description of this method was given by Archimedes. Essentially, the method is a limiting process in which the area is squeezed between two polygons—one inscribed in the region and one circumscribed about the region.

For instance, in Figure 4.8 the area of a circular region is approximated by an  $n$ -sided inscribed polygon and an  $n$ -sided circumscribed polygon. For each value of  $n$  the area of the inscribed polygon is less than the area of the circle, and the area of the circumscribed polygon is greater than the area of the circle. Moreover, as  $n$  increases, the areas of both polygons become better and better approximations of the area of the circle.



The exhaustion method for finding the area of a circular region  
Figure 4.8

A process that is similar to that used by Archimedes to determine the area of a plane region is used in the remaining examples in this section.



Mary Evans Picture Library

ARCHIMEDES (287–212 B.C.)

Archimedes used the method of exhaustion to derive formulas for the areas of ellipses, parabolic segments, and sectors of a spiral. He is considered to have been the greatest applied mathematician of antiquity.

**FOR FURTHER INFORMATION** For an alternative development of the formula for the area of a circle, see the article “Proof Without Words: Area of a Disk is  $\pi R^2$ ” by Russell Jay Hendel in *Mathematics Magazine*. To view this article, go to the website [www.matharticles.com](http://www.matharticles.com).

## The Area of a Plane Region

Recall from Section 1.1 that the origins of calculus are connected to two classic problems: the tangent line problem and the area problem. Example 3 begins the investigation of the area problem.

### EXAMPLE 3 Approximating the Area of a Plane Region

Use the five rectangles in Figure 4.9(a) and (b) to find *two* approximations of the area of the region lying between the graph of

$$f(x) = -x^2 + 5$$

and the  $x$ -axis between  $x = 0$  and  $x = 2$ .

#### Solution

- a.** The right endpoints of the five intervals are  $\frac{2}{5}i$ , where  $i = 1, 2, 3, 4, 5$ . The width of each rectangle is  $\frac{2}{5}$ , and the height of each rectangle can be obtained by evaluating  $f$  at the right endpoint of each interval.

$$\left[0, \frac{2}{5}\right], \left[\frac{2}{5}, \frac{4}{5}\right], \left[\frac{4}{5}, \frac{6}{5}\right], \left[\frac{6}{5}, \frac{8}{5}\right], \left[\frac{8}{5}, \frac{10}{5}\right]$$



Evaluate  $f$  at the right endpoints of these intervals.

The sum of the areas of the five rectangles is

$$\sum_{i=1}^5 \overbrace{f\left(\frac{2i}{5}\right)}^{\text{Height}} \overbrace{\left(\frac{2}{5}\right)}^{\text{Width}} = \sum_{i=1}^5 \left[ -\left(\frac{2i}{5}\right)^2 + 5 \right] \left(\frac{2}{5}\right) = \frac{162}{25} = 6.48.$$

Because each of the five rectangles lies inside the parabolic region, you can conclude that the area of the parabolic region is greater than 6.48.

- b.** The left endpoints of the five intervals are  $\frac{2}{5}(i - 1)$ , where  $i = 1, 2, 3, 4, 5$ . The width of each rectangle is  $\frac{2}{5}$ , and the height of each rectangle can be obtained by evaluating  $f$  at the left endpoint of each interval.

$$\sum_{i=1}^5 \overbrace{f\left(\frac{2i-2}{5}\right)}^{\text{Height}} \overbrace{\left(\frac{2}{5}\right)}^{\text{Width}} = \sum_{i=1}^5 \left[ -\left(\frac{2i-2}{5}\right)^2 + 5 \right] \left(\frac{2}{5}\right) = \frac{202}{25} = 8.08.$$


Because the parabolic region lies within the union of the five rectangular regions, you can conclude that the area of the parabolic region is less than 8.08.

By combining the results in parts (a) and (b), you can conclude that


$$6.48 < (\text{Area of region}) < 8.08.$$

**NOTE** By increasing the number of rectangles used in Example 3, you can obtain closer and closer approximations of the area of the region. For instance, using 25 rectangles of width  $\frac{2}{25}$  each, you can conclude that

$$7.17 < (\text{Area of region}) < 7.49.$$




(a) The area of the parabolic region is greater than the area of the rectangles.




(b) The area of the parabolic region is less than the area of the rectangles.

**Figure 4.9**





The region under a curve  
Figure 4.10



The interval  $[a, b]$  is divided into  $n$   
subintervals of width  $\Delta x = \frac{b-a}{n}$ .

Figure 4.11

## Upper and Lower Sums

The procedure used in Example 3 can be generalized as follows. Consider a plane region bounded above by the graph of a nonnegative, continuous function  $y = f(x)$ , as shown in Figure 4.10. The region is bounded below by the  $x$ -axis, and the left and right boundaries of the region are the vertical lines  $x = a$  and  $x = b$ .

To approximate the area of the region, begin by subdividing the interval  $[a, b]$  into  $n$  subintervals, each of width  $\Delta x = (b - a)/n$ , as shown in Figure 4.11. The endpoints of the intervals are as follows.

$$\begin{array}{ccccccc} a = x_0 & & x_1 & & x_2 & & x_n = b \\ a + 0(\Delta x) & < & a + 1(\Delta x) & < & a + 2(\Delta x) & < \cdots < & a + n(\Delta x) \end{array}$$

Because  $f$  is continuous, the Extreme Value Theorem guarantees the existence of a minimum and a maximum value of  $f(x)$  in *each* subinterval.

$f(m_i)$  = Minimum value of  $f(x)$  in  $i$ th subinterval

$f(M_i)$  = Maximum value of  $f(x)$  in  $i$ th subinterval

Next, define an **inscribed rectangle** lying *inside* the  $i$ th subregion and a **circumscribed rectangle** extending *outside* the  $i$ th subregion. The height of the  $i$ th inscribed rectangle is  $f(m_i)$  and the height of the  $i$ th circumscribed rectangle is  $f(M_i)$ . For *each*  $i$ , the area of the inscribed rectangle is less than or equal to the area of the circumscribed rectangle.

$$\left( \begin{array}{c} \text{Area of inscribed} \\ \text{rectangle} \end{array} \right) = f(m_i) \Delta x \leq f(M_i) \Delta x = \left( \begin{array}{c} \text{Area of circumscribed} \\ \text{rectangle} \end{array} \right)$$

The sum of the areas of the inscribed rectangles is called a **lower sum**, and the sum of the areas of the circumscribed rectangles is called an **upper sum**.

$$\text{Lower sum} = s(n) = \sum_{i=1}^n f(m_i) \Delta x \quad \text{Area of inscribed rectangles}$$

$$\text{Upper sum} = S(n) = \sum_{i=1}^n f(M_i) \Delta x \quad \text{Area of circumscribed rectangles}$$

From Figure 4.12, you can see that the lower sum  $s(n)$  is less than or equal to the upper sum  $S(n)$ . Moreover, the actual area of the region lies between these two sums.

$$s(n) \leq (\text{Area of region}) \leq S(n)$$






Figure 4.12



Inscribed rectangles



Circumscribed rectangles

Figure 4.13

**EXAMPLE 4** Finding Upper and Lower Sums for a Region

Find the upper and lower sums for the region bounded by the graph of  $f(x) = x^2$  and the  $x$ -axis between  $x = 0$  and  $x = 2$ .

**Solution** To begin, partition the interval  $[0, 2]$  into  $n$  subintervals, each of width

$$\Delta x = \frac{b - a}{n} = \frac{2 - 0}{n} = \frac{2}{n}.$$

Figure 4.13 shows the endpoints of the subintervals and several inscribed and circumscribed rectangles. Because  $f$  is increasing on the interval  $[0, 2]$ , the minimum value on each subinterval occurs at the left endpoint, and the maximum value occurs at the right endpoint.

Left Endpoints

$$m_i = 0 + (i - 1)\left(\frac{2}{n}\right) = \frac{2(i - 1)}{n}$$

Right Endpoints

$$M_i = 0 + i\left(\frac{2}{n}\right) = \frac{2i}{n}$$

Using the left endpoints, the lower sum is

$$\begin{aligned} s(n) &= \sum_{i=1}^n f(m_i) \Delta x = \sum_{i=1}^n f\left[\frac{2(i-1)}{n}\right] \left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left[\frac{2(i-1)}{n}\right]^2 \left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left(\frac{8}{n^3}\right)(i^2 - 2i + 1) \\ &= \frac{8}{n^3} \left( \sum_{i=1}^n i^2 - 2 \sum_{i=1}^n i + \sum_{i=1}^n 1 \right) \\ &= \frac{8}{n^3} \left\{ \frac{n(n+1)(2n+1)}{6} - 2 \left[ \frac{n(n+1)}{2} \right] + n \right\} \\ &= \frac{4}{3n^3} (2n^3 - 3n^2 + n) \\ &= \frac{8}{3} - \frac{4}{n} + \frac{4}{3n^2}. \end{aligned}$$

Lower sum

Using the right endpoints, the upper sum is

$$\begin{aligned} S(n) &= \sum_{i=1}^n f(M_i) \Delta x = \sum_{i=1}^n f\left(\frac{2i}{n}\right) \left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left(\frac{2i}{n}\right)^2 \left(\frac{2}{n}\right) \\ &= \sum_{i=1}^n \left(\frac{8}{n^3}\right) i^2 \\ &= \frac{8}{n^3} \left[ \frac{n(n+1)(2n+1)}{6} \right] \\ &= \frac{4}{3n^3} (2n^3 + 3n^2 + n) \\ &= \frac{8}{3} + \frac{4}{n} + \frac{4}{3n^2}. \end{aligned}$$

Upper sum

**EXPLORATION**

For the region given in Example 4, evaluate the lower sum

$$s(n) = \frac{8}{3} - \frac{4}{n} + \frac{4}{3n^2}$$

and the upper sum

$$S(n) = \frac{8}{3} + \frac{4}{n} + \frac{4}{3n^2}$$

for  $n = 10, 100$ , and  $1000$ . Use your results to determine the area of the region.

Example 4 illustrates some important things about lower and upper sums. First, notice that for any value of  $n$ , the lower sum is less than (or equal to) the upper sum.

$$s(n) = \frac{8}{3} - \frac{4}{n} + \frac{4}{3n^2} < \frac{8}{3} + \frac{4}{n} + \frac{4}{3n^2} = S(n)$$

Second, the difference between these two sums lessens as  $n$  increases. In fact, if you take the limits as  $n \rightarrow \infty$ , both the upper sum and the lower sum approach  $\frac{8}{3}$ .

$$\lim_{n \rightarrow \infty} s(n) = \lim_{n \rightarrow \infty} \left( \frac{8}{3} - \frac{4}{n} + \frac{4}{3n^2} \right) = \frac{8}{3} \quad \text{Lower sum limit}$$

$$\lim_{n \rightarrow \infty} S(n) = \lim_{n \rightarrow \infty} \left( \frac{8}{3} + \frac{4}{n} + \frac{4}{3n^2} \right) = \frac{8}{3} \quad \text{Upper sum limit}$$

The next theorem shows that the equivalence of the limits (as  $n \rightarrow \infty$ ) of the upper and lower sums is not mere coincidence. It is true for all functions that are continuous and nonnegative on the closed interval  $[a, b]$ . The proof of this theorem is best left to a course in advanced calculus.

**THEOREM 4.3 Limits of the Lower and Upper Sums**

Let  $f$  be continuous and nonnegative on the interval  $[a, b]$ . The limits as  $n \rightarrow \infty$  of both the lower and upper sums exist and are equal to each other. That is,

$$\begin{aligned} \lim_{n \rightarrow \infty} s(n) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(m_i) \Delta x \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(M_i) \Delta x \\ &= \lim_{n \rightarrow \infty} S(n) \end{aligned}$$

where  $\Delta x = (b - a)/n$  and  $f(m_i)$  and  $f(M_i)$  are the minimum and maximum values of  $f$  on the subinterval.


Because the same limit is attained for both the minimum value  $f(m_i)$  and the maximum value  $f(M_i)$ , it follows from the Squeeze Theorem (Theorem 1.8) that the choice of  $x$  in the  $i$ th subinterval does not affect the limit. This means that you are free to choose an *arbitrary*  $x$ -value in the  $i$ th subinterval, as in the following *definition of the area of a region in the plane*.

**Definition of the Area of a Region in the Plane**

Let  $f$  be continuous and nonnegative on the interval  $[a, b]$ . The area of the region bounded by the graph of  $f$ , the  $x$ -axis, and the vertical lines  $x = a$  and  $x = b$  is

$$\text{Area} = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x, \quad x_{i-1} \leq c_i \leq x_i$$





where  $\Delta x = (b - a)/n$  (see Figure 4.14).



The width of the  $i$ th subinterval is  $\Delta x = x_i - x_{i-1}$ .


Figure 4.14

In Exercises 23–26, bound the area of the shaded region by approximating the upper and lower sums. Use rectangles of width 1.




In Exercises 27–30, use upper and lower sums to approximate the area of the region using the given number of subintervals (of equal width).


27.  $y = \sqrt{x}$




28.  $y = \sqrt{x} + 2$



29.  $y = \frac{1}{x}$



30.  $y = \sqrt{1 - x^2}$



In Exercises 31–34, find the limit of  $s(n)$  as  $n \rightarrow \infty$ .

31.  $s(n) = \frac{81}{n^4} \left[ \frac{n^2(n+1)^2}{4} \right]$

32.  $s(n) = \frac{64}{n^3} \left[ \frac{n(n+1)(2n+1)}{6} \right]$

33.  $s(n) = \frac{18}{n^2} \left[ \frac{n(n+1)}{2} \right]$

34.  $s(n) = \frac{1}{n^2} \left[ \frac{n(n+1)}{2} \right]$

In Exercises 35–38, use the summation formulas to rewrite the expression without the summation notation. Use the result to find the sum for  $n = 10, 100, 1000$ , and  $10,000$ .

35.  $\sum_{i=1}^n \frac{2i+1}{n^2}$

36.  $\sum_{j=1}^n \frac{4j+3}{n^2}$

37.  $\sum_{k=1}^n \frac{6k(k-1)}{n^3}$

38.  $\sum_{i=1}^n \frac{4i^2(i-1)}{n^4}$

In Exercises 39–44, find a formula for the sum of  $n$  terms. Use the formula to find the limit as  $n \rightarrow \infty$ .

39.  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{16i}{n^2}$

40.  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \left( \frac{2i}{n} \right) \left( \frac{2}{n} \right)$

41.  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n^3} (i-1)^2$

42.  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \left( 1 + \frac{2i}{n} \right)^2 \left( \frac{2}{n} \right)$

43.  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \left( 1 + \frac{i}{n} \right) \left( \frac{2}{n} \right)$

44.  $\lim_{n \rightarrow \infty} \sum_{i=1}^n \left( 1 + \frac{2i}{n} \right)^3 \left( \frac{2}{n} \right)$

45. **Numerical Reasoning** Consider a triangle of area 2 bounded by the graphs of  $y = x$ ,  $y = 0$ , and  $x = 2$ .

(a) Sketch the region.

(b) Divide the interval  $[0, 2]$  into  $n$  subintervals of equal width and show that the endpoints are

$$0 < 1\left(\frac{2}{n}\right) < \cdots < (n-1)\left(\frac{2}{n}\right) < n\left(\frac{2}{n}\right).$$

(c) Show that  $s(n) = \sum_{i=1}^n \left[ (i-1)\left(\frac{2}{n}\right) \right] \left( \frac{2}{n} \right)$ .

(d) Show that  $S(n) = \sum_{i=1}^n \left[ i\left(\frac{2}{n}\right) \right] \left( \frac{2}{n} \right)$ .

(e) Complete the table.

$n$	5	10	50	100
$s(n)$				
$S(n)$				

(f) Show that  $\lim_{n \rightarrow \infty} s(n) = \lim_{n \rightarrow \infty} S(n) = 2$ .

46. **Numerical Reasoning** Consider a trapezoid of area 4 bounded by the graphs of  $y = x$ ,  $y = 0$ ,  $x = 1$ , and  $x = 3$ .

(a) Sketch the region.

(b) Divide the interval  $[1, 3]$  into  $n$  subintervals of equal width and show that the endpoints are

$$1 < 1 + 1\left(\frac{2}{n}\right) < \cdots < 1 + (n-1)\left(\frac{2}{n}\right) < 1 + n\left(\frac{2}{n}\right).$$

(c) Show that  $s(n) = \sum_{i=1}^n \left[ 1 + (i-1)\left(\frac{2}{n}\right) \right] \left( \frac{2}{n} \right)$ .

(d) Show that  $S(n) = \sum_{i=1}^n \left[ 1 + i\left(\frac{2}{n}\right) \right] \left( \frac{2}{n} \right)$ .

(e) Complete the table.

$n$	5	10	50	100
$s(n)$				
$S(n)$				

(f) Show that  $\lim_{n \rightarrow \infty} s(n) = \lim_{n \rightarrow \infty} S(n) = 4$ .

In Exercises 47–56, use the limit process to find the area of the region between the graph of the function and the  $x$ -axis over the given interval. Sketch the region.

47.  $y = -2x + 3$ ,  $[0, 1]$       48.  $y = 3x - 4$ ,  $[2, 5]$   
 49.  $y = x^2 + 2$ ,  $[0, 1]$       50.  $y = x^2 + 1$ ,  $[0, 3]$   
 51.  $y = 16 - x^2$ ,  $[1, 3]$       52.  $y = 1 - x^2$ ,  $[-1, 1]$   
 53.  $y = 64 - x^3$ ,  $[1, 4]$       54.  $y = 2x - x^3$ ,  $[0, 1]$   
 55.  $y = x^2 - x^3$ ,  $[-1, 1]$       56.  $y = x^2 - x^3$ ,  $[-1, 0]$

In Exercises 57–62, use the limit process to find the area of the region between the graph of the function and the  $y$ -axis over the given  $y$ -interval. Sketch the region.


57.  $f(y) = 3y$ ,  $0 \leq y \leq 2$       58.  $g(y) = \frac{1}{2}y$ ,  $2 \leq y \leq 4$   
 59.  $f(y) = y^2$ ,  $0 \leq y \leq 3$       60.  $f(y) = 4y - y^2$ ,  $1 \leq y \leq 2$   
 61.  $g(y) = 4y^2 - y^3$ ,  $1 \leq y \leq 3$       62.  $h(y) = y^3 + 1$ ,  $1 \leq y \leq 2$

In Exercises 63–66, use the *Midpoint Rule*

$$\text{Area} \approx \sum_{i=1}^n f\left(\frac{x_i + x_{i-1}}{2}\right) \Delta x$$

with  $n = 4$  to approximate the area of the region bounded by the graph of the function and the  $x$ -axis over the given interval.

63.  $f(x) = x^2 + 3$ ,  $[0, 2]$       64.  $f(x) = x^2 + 4x$ ,  $[0, 4]$   
 65.  $f(x) = \tan x$ ,  $\left[0, \frac{\pi}{4}\right]$       66.  $f(x) = \sin x$ ,  $\left[0, \frac{\pi}{2}\right]$

 **Programming** Write a program for a graphing utility to approximate areas by using the Midpoint Rule. Assume that the function is positive over the given interval and the subintervals are of equal width. In Exercises 67–70, use the program to approximate the area of the region between the graph of the function and the  $x$ -axis over the given interval, and complete the table.

$n$	4	8	12	16	20
Approximate Area					

67.  $f(x) = \sqrt{x}$ ,  $[0, 4]$       68.  $f(x) = \frac{8}{x^2 + 1}$ ,  $[2, 6]$   
 69.  $f(x) = \tan\left(\frac{\pi x}{8}\right)$ ,  $[1, 3]$       70.  $f(x) = \cos \sqrt{x}$ ,  $[0, 2]$

### Writing About Concepts

**Approximation** In Exercises 71 and 72, determine which value best approximates the area of the region between the  $x$ -axis and the graph of the function over the given interval. (Make your selection on the basis of a sketch of the region and not by performing calculations.)

71.  $f(x) = 4 - x^2$ ,  $[0, 2]$   
 (a)  $-2$    (b)  $6$    (c)  $10$    (d)  $3$    (e)  $8$


### Writing About Concepts (continued)

72.  $f(x) = \sin \frac{\pi x}{4}$ ,  $[0, 4]$   
 (a)  $3$    (b)  $1$    (c)  $-2$    (d)  $8$    (e)  $6$
73. In your own words and using appropriate figures, describe the methods of upper sums and lower sums in approximating the area of a region.
74. Give the definition of the area of a region in the plane.

**75. Graphical Reasoning** Consider the region bounded by the graphs of

$$f(x) = \frac{8x}{x+1},$$

$x = 0$ ,  $x = 4$ , and  $y = 0$ , as shown in the figure. To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).




- (a) Redraw the figure, and complete and shade the rectangles representing the lower sum when  $n = 4$ . Find this lower sum.
- (b) Redraw the figure, and complete and shade the rectangles representing the upper sum when  $n = 4$ . Find this upper sum.
- (c) Redraw the figure, and complete and shade the rectangles whose heights are determined by the functional values at the midpoint of each subinterval when  $n = 4$ . Find this sum using the Midpoint Rule.
- (d) Verify the following formulas for approximating the area of the region using  $n$  subintervals of equal width.

$$\text{Lower sum: } s(n) = \sum_{i=1}^n f\left[\left(i-1\right)\frac{4}{n}\right]\left(\frac{4}{n}\right)$$

$$\text{Upper sum: } S(n) = \sum_{i=1}^n f\left[\left(i\right)\frac{4}{n}\right]\left(\frac{4}{n}\right)$$

$$\text{Midpoint Rule: } M(n) = \sum_{i=1}^n f\left[\left(i-\frac{1}{2}\right)\frac{4}{n}\right]\left(\frac{4}{n}\right)$$

-  (e) Use a graphing utility and the formulas in part (d) to complete the table.

$n$	4	8	20	100	200
$s(n)$					
$S(n)$					
$M(n)$					

- (f) Explain why  $s(n)$  increases and  $S(n)$  decreases for increasing values of  $n$ , as shown in the table in part (e).

**76. Monte Carlo Method** The following computer program approximates the area of the region under the graph of a monotonic function and above the  $x$ -axis between  $x = a$  and  $x = b$ . Run the program for  $a = 0$  and  $b = \pi/2$  for several values of  $N2$ . Explain why the Monte Carlo Method works. [Adaptation of Monte Carlo Method program from James M. Sconyers, "Approximation of Area Under a Curve," *MATHEMATICS TEACHER* 77, no. 2 (February 1984). Copyright © 1984 by the National Council of Teachers of Mathematics. Reprinted with permission.]

```

10 DEF FNF(X)=SIN(X)
20 A=0
30 B=PI/2
40 PRINT "Input Number of Random Points"
50 INPUT N2
60 N1=0
70 IF FNF(A)>FNF(B) THEN YMAX=FNF(A) ELSE
   YMAX=FNF(B)
80 FOR I=1 TO N2
90 X=A+(B-A)*RND(1)
100 Y=YMAX*RND(1)
110 IF Y>=FNF(X) THEN GOTO 130
120 N1=N1+1
130 NEXT I
140 AREA=(N1/N2)*(B-A)*YMAX
150 PRINT "Approximate Area:"; AREA
160 END

```

**True or False?** In Exercises 77 and 78, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

77. The sum of the first  $n$  positive integers is  $n(n+1)/2$ .
78. If  $f$  is continuous and nonnegative on  $[a, b]$ , then the limits as  $n \rightarrow \infty$  of its lower sum  $s(n)$  and upper sum  $S(n)$  both exist and are equal.
79. **Writing** Use the figure to write a short paragraph explaining why the formula  $1 + 2 + \cdots + n = \frac{1}{2}n(n+1)$  is valid for all positive integers  $n$ .



Figure 79




Figure 80

**80. Graphical Reasoning** Consider an  $n$ -sided regular polygon inscribed in a circle of radius  $r$ . Join the vertices of the polygon to the center of the circle, forming  $n$  congruent triangles (see figure).

- (a) Determine the central angle  $\theta$  in terms of  $n$ .
- (b) Show that the area of each triangle is  $\frac{1}{2}r^2 \sin \theta$ .
- (c) Let  $A_n$  be the sum of the areas of the  $n$  triangles. Find  $\lim_{n \rightarrow \infty} A_n$ .



**81. Modeling Data** The table lists the measurements of a lot bounded by a stream and two straight roads that meet at right angles, where  $x$  and  $y$  are measured in feet (see figure).

$x$	0	50	100	150	200	250	300
$y$	450	362	305	268	245	156	0

- (a) Use the regression capabilities of a graphing utility to find a model of the form  $y = ax^3 + bx^2 + cx + d$ .
- (b) Use a graphing utility to plot the data and graph the model.
- (c) Use the model in part (a) to estimate the area of the lot.




Figure for 81




Figure for 82

**82. Building Blocks** A child places  $n$  cubic building blocks in a row to form the base of a triangular design (see figure). Each successive row contains two fewer blocks than the preceding row. Find a formula for the number of blocks used in the design. (Hint: The number of building blocks in the design depends on whether  $n$  is odd or even.)

**83.** Prove each formula by mathematical induction. (You may need to review the method of proof by induction from a precalculus text.)

(a)  $\sum_{i=1}^n 2i = n(n+1)$

(b)  $\sum_{i=1}^n i^3 = \frac{n^2(n+1)^2}{4}$

### Putnam Exam Challenge

- 84.** A dart, thrown at random, hits a square target. Assuming that any two parts of the target of equal area are equally likely to be hit, find the probability that the point hit is nearer to the center than to any edge. Write your answer in the form  $(a\sqrt{b} + c)/d$ , where  $a$ ,  $b$ ,  $c$ , and  $d$  are positive integers.

This problem was composed by the Committee on the Putnam Prize Competition.  
© The Mathematical Association of America. All rights reserved.



## Section 4.3

## Riemann Sums and Definite Integrals

- Understand the definition of a Riemann sum.
- Evaluate a definite integral using limits.
- Evaluate a definite integral using properties of definite integrals.

## Riemann Sums

In the definition of area given in Section 4.2, the partitions have subintervals of *equal width*. This was done only for computational convenience. The following example shows that it is not necessary to have subintervals of equal width.

**EXAMPLE 1** A Partition with Subintervals of Unequal Widths

Consider the region bounded by the graph of  $f(x) = \sqrt{x}$  and the  $x$ -axis for  $0 \leq x \leq 1$ , as shown in Figure 4.18. Evaluate the limit

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i$$

where  $c_i$  is the right endpoint of the partition given by  $c_i = i^2/n^2$  and  $\Delta x_i$  is the width of the  $i$ th interval.


**Solution** The width of the  $i$ th interval is given by

$$\begin{aligned} \Delta x_i &= \frac{i^2}{n^2} - \frac{(i-1)^2}{n^2} \\ &= \frac{i^2 - i^2 + 2i - 1}{n^2} \\ &= \frac{2i - 1}{n^2}. \end{aligned}$$


So, the limit is

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{\frac{i^2}{n^2}} \left( \frac{2i - 1}{n^2} \right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{i=1}^n (2i^2 - i) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^3} \left[ 2 \left( \frac{n(n+1)(2n+1)}{6} \right) - \frac{n(n+1)}{2} \right] \\ &= \lim_{n \rightarrow \infty} \frac{4n^3 + 3n^2 - n}{6n^3} \\ &= \frac{2}{3}. \end{aligned}$$

From Example 7 in Section 4.2, you know that the region shown in Figure 4.19 has an area of  $\frac{1}{3}$ . Because the square bounded by  $0 \leq x \leq 1$  and  $0 \leq y \leq 1$  has an area of 1, you can conclude that the area of the region shown in Figure 4.18 has an area of  $\frac{2}{3}$ . This agrees with the limit found in Example 1, even though that example used a partition having subintervals of unequal widths. The reason this particular partition gave the proper area is that as  $n$  increases, the *width of the largest subinterval approaches zero*. This is a key feature of the development of definite integrals.



The subintervals do not have equal widths.  
**Figure 4.18**



The area of the region bounded by the graph of  $x = y^2$  and the  $y$ -axis for  $0 \leq y \leq 1$  is  $\frac{1}{3}$ .  
**Figure 4.19**



In the preceding section, the limit of a sum was used to define the area of a region in the plane. Finding area by this means is only one of *many* applications involving the limit of a sum. A similar approach can be used to determine quantities as diverse as arc lengths, average values, centroids, volumes, work, and surface areas. The following definition is named after Georg Friedrich Bernhard Riemann. Although the definite integral had been defined and used long before the time of Riemann, he generalized the concept to cover a broader category of functions.

In the following definition of a Riemann sum, note that the function  $f$  has no restrictions other than being defined on the interval  $[a, b]$ . (In the preceding section, the function  $f$  was assumed to be continuous and nonnegative because we were dealing with the area under a curve.)

### Definition of a Riemann Sum

Let  $f$  be defined on the closed interval  $[a, b]$ , and let  $\Delta$  be a partition of  $[a, b]$  given by

$$a = x_0 < x_1 < x_2 < \cdots < x_{n-1} < x_n = b$$

where  $\Delta x_i$  is the width of the  $i$ th subinterval. If  $c_i$  is any point in the  $i$ th subinterval, then the sum

$$\sum_{i=1}^n f(c_i) \Delta x_i, \quad x_{i-1} \leq c_i \leq x_i$$

is called a **Riemann sum** of  $f$  for the partition  $\Delta$ .

**NOTE** The sums in Section 4.2 are examples of Riemann sums, but there are more general Riemann sums than those covered there.

The width of the largest subinterval of a partition  $\Delta$  is the **norm** of the partition and is denoted by  $\|\Delta\|$ . If every subinterval is of equal width, the partition is **regular** and the norm is denoted by

$$\|\Delta\| = \Delta x = \frac{b-a}{n}, \quad \text{Regular partition}$$

For a general partition, the norm is related to the number of subintervals of  $[a, b]$  in the following way.


$$\frac{b-a}{\|\Delta\|} \leq n \quad \text{General partition}$$

So, the number of subintervals in a partition approaches infinity as the norm of the partition approaches 0. That is,  $\|\Delta\| \rightarrow 0$  implies that  $n \rightarrow \infty$ .

The converse of this statement is not true. For example, let  $\Delta_n$  be the partition of the interval  $[0, 1]$  given by

$$0 < \frac{1}{2^n} < \frac{1}{2^{n-1}} < \cdots < \frac{1}{8} < \frac{1}{4} < \frac{1}{2} < 1.$$

As shown in Figure 4.20, for any positive value of  $n$ , the norm of the partition  $\Delta_n$  is  $\frac{1}{2}$ . So, letting  $n$  approach infinity does not force  $\|\Delta\|$  to approach 0. In a regular partition, however, the statements  $\|\Delta\| \rightarrow 0$  and  $n \rightarrow \infty$  are equivalent.



$n \rightarrow \infty$  does not imply that  $\|\Delta\| \rightarrow 0$ .  
**Figure 4.20**

## Definite Integrals

To define the definite integral, consider the following limit.

$$\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n f(c_i) \Delta x_i = L$$

To say that this limit exists means that for  $\varepsilon > 0$  there exists a  $\delta > 0$  such that for every partition with  $\|\Delta\| < \delta$  it follows that

$$\left| L - \sum_{i=1}^n f(c_i) \Delta x_i \right| < \varepsilon.$$

(This must be true for any choice of  $c_i$  in the  $i$ th subinterval of  $\Delta$ .)

**FOR FURTHER INFORMATION** For insight into the history of the definite integral, see the article “The Evolution of Integration” by A. Shenitzer and J. Steprāns in *The American Mathematical Monthly*. To view this article, go to the website [www.matharticles.com](http://www.matharticles.com).

### Definition of a Definite Integral

If  $f$  is defined on the closed interval  $[a, b]$  and the limit

$$\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n f(c_i) \Delta x_i$$

exists (as described above), then  $f$  is **integrable** on  $[a, b]$  and the limit is denoted by

$$\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n f(c_i) \Delta x_i = \int_a^b f(x) dx.$$

The limit is called the **definite integral** of  $f$  from  $a$  to  $b$ . The number  $a$  is the **lower limit** of integration, and the number  $b$  is the **upper limit** of integration.

It is not a coincidence that the notation for definite integrals is similar to that used for indefinite integrals. You will see why in the next section when the Fundamental Theorem of Calculus is introduced. For now it is important to see that definite integrals and indefinite integrals are different identities. A definite integral is a *number*, whereas an indefinite integral is a *family of functions*.

A sufficient condition for a function  $f$  to be integrable on  $[a, b]$  is that it is continuous on  $[a, b]$ . A proof of this theorem is beyond the scope of this text.

### THEOREM 4.4 Continuity Implies Integrability

If a function  $f$  is continuous on the closed interval  $[a, b]$ , then  $f$  is integrable on  $[a, b]$ .

### EXPLORATION

**The Converse of Theorem 4.4** Is the converse of Theorem 4.4 true? That is, if a function is integrable, does it have to be continuous? Explain your reasoning and give examples.

Describe the relationships among continuity, differentiability, and integrability. Which is the strongest condition? Which is the weakest? Which conditions imply other conditions?

**EXAMPLE 2** Evaluating a Definite Integral as a Limit

Evaluate the definite integral  $\int_{-2}^1 2x \, dx$ .

**Solution** The function  $f(x) = 2x$  is integrable on the interval  $[-2, 1]$  because it is continuous on  $[-2, 1]$ . Moreover, the definition of integrability implies that any partition whose norm approaches 0 can be used to determine the limit. For computational convenience, define  $\Delta$  by subdividing  $[-2, 1]$  into  $n$  subintervals of equal width


$$\Delta x_i = \Delta x = \frac{b - a}{n} = \frac{3}{n}.$$

Choosing  $c_i$  as the right endpoint of each subinterval produces

$$c_i = a + i(\Delta x) = -2 + \frac{3i}{n}.$$

So, the definite integral is given by

$$\begin{aligned} \int_{-2}^1 2x \, dx &= \lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n f(c_i) \Delta x_i \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n 2 \left( -2 + \frac{3i}{n} \right) \left( \frac{3}{n} \right) \\ &= \lim_{n \rightarrow \infty} \frac{6}{n} \sum_{i=1}^n \left( -2 + \frac{3i}{n} \right) \\ &= \lim_{n \rightarrow \infty} \frac{6}{n} \left\{ -2n + \frac{3}{n} \left[ \frac{n(n+1)}{2} \right] \right\} \\ &= \lim_{n \rightarrow \infty} \left( -12 + 9 + \frac{9}{n} \right) \\ &= -3. \end{aligned}$$



Because the definite integral is negative, it does not represent the area of the region.

**Figure 4.21**


Because the definite integral in Example 2 is negative, it *does not* represent the area of the region shown in Figure 4.21. Definite integrals can be positive, negative, or zero. For a definite integral to be interpreted as an area (as defined in Section 4.2), the function  $f$  must be continuous and nonnegative on  $[a, b]$ , as stated in the following theorem. (The proof of this theorem is straightforward—you simply use the definition of area given in Section 4.2.)

**THEOREM 4.5** The Definite Integral as the Area of a Region

If  $f$  is continuous and nonnegative on the closed interval  $[a, b]$ , then the area of the region bounded by the graph of  $f$ , the  $x$ -axis, and the vertical lines  $x = a$  and  $x = b$  is given by


$$\text{Area} = \int_a^b f(x) \, dx.$$

(See Figure 4.22.)



You can use a definite integral to find the area of the region bounded by the graph of  $f$ , the  $x$ -axis,  $x = a$ , and  $x = b$ .

**Figure 4.22**



$$\text{Area} = \int_0^4 (4x - x^2) dx$$

Figure 4.23

As an example of Theorem 4.5, consider the region bounded by the graph of

$$f(x) = 4x - x^2$$

and the  $x$ -axis, as shown in Figure 4.23. Because  $f$  is continuous and nonnegative on the closed interval  $[0, 4]$ , the area of the region is

$$\text{Area} = \int_0^4 (4x - x^2) dx.$$

A straightforward technique for evaluating a definite integral such as this will be discussed in Section 4.4. For now, however, you can evaluate a definite integral in two ways—you can use the limit definition *or* you can check to see whether the definite integral represents the area of a common geometric region such as a rectangle, triangle, or semicircle.

### EXAMPLE 3 Areas of Common Geometric Figures

Sketch the region corresponding to each definite integral. Then evaluate each integral using a geometric formula.

a.  $\int_1^3 4 dx$       b.  $\int_0^3 (x + 2) dx$       c.  $\int_{-2}^2 \sqrt{4 - x^2} dx$

**Solution** A sketch of each region is shown in Figure 4.24.

a. This region is a rectangle of height 4 and width 2.

$$\int_1^3 4 dx = (\text{Area of rectangle}) = 4(2) = 8$$

b. This region is a trapezoid with an altitude of 3 and parallel bases of lengths 2 and 5. The formula for the area of a trapezoid is  $\frac{1}{2}h(b_1 + b_2)$ .

$$\int_0^3 (x + 2) dx = (\text{Area of trapezoid}) = \frac{1}{2}(3)(2 + 5) = \frac{21}{2}$$

c. This region is a semicircle of radius 2. The formula for the area of a semicircle is  $\frac{1}{2}\pi r^2$ .

$$\int_{-2}^2 \sqrt{4 - x^2} dx = (\text{Area of semicircle}) = \frac{1}{2}\pi(2^2) = 2\pi$$


**NOTE** The variable of integration in a definite integral is sometimes called a *dummy variable* because it can be replaced by any other variable without changing the value of the integral. For instance, the definite integrals

$$\int_0^3 (x + 2) dx$$


and

$$\int_0^3 (t + 2) dt$$


have the same value.



(a)



(b)



(c)

Figure 4.24

## Properties of Definite Integrals

The definition of the definite integral of  $f$  on the interval  $[a, b]$  specifies that  $a < b$ . Now, however, it is convenient to extend the definition to cover cases in which  $a = b$  or  $a > b$ . Geometrically, the following two definitions seem reasonable. For instance, it makes sense to define the area of a region of zero width and finite height to be 0.

### Definitions of Two Special Definite Integrals

1. If  $f$  is defined at  $x = a$ , then we define  $\int_a^a f(x) dx = 0$ .
2. If  $f$  is integrable on  $[a, b]$ , then we define  $\int_b^a f(x) dx = -\int_a^b f(x) dx$ .



### EXAMPLE 4 Evaluating Definite Integrals

- a. Because the sine function is defined at  $x = \pi$ , and the upper and lower limits of integration are equal, you can write

$$\int_{\pi}^{\pi} \sin x dx = 0.$$

- b. The integral  $\int_3^0 (x + 2) dx$  is the same as that given in Example 3(b) except that the upper and lower limits are interchanged. Because the integral in Example 3(b) has a value of  $\frac{21}{2}$ , you can write

$$\int_3^0 (x + 2) dx = -\int_0^3 (x + 2) dx = -\frac{21}{2}.$$

In Figure 4.25, the larger region can be divided at  $x = c$  into two subregions whose intersection is a line segment. Because the line segment has zero area, it follows that the area of the larger region is equal to the sum of the areas of the two smaller regions.




Figure 4.25

### THEOREM 4.6 Additive Interval Property

If  $f$  is integrable on the three closed intervals determined by  $a$ ,  $b$ , and  $c$ , then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

### EXAMPLE 5 Using the Additive Interval Property

$$\begin{aligned} \int_{-1}^1 |x| dx &= \int_{-1}^0 -x dx + \int_0^1 x dx && \text{Theorem 4.6} \\ &= \frac{1}{2} + \frac{1}{2} && \text{Area of a triangle} \\ &= 1 \end{aligned}$$



Because the definite integral is defined as the limit of a sum, it inherits the properties of summation given at the top of page 260.

### THEOREM 4.7 Properties of Definite Integrals

If  $f$  and  $g$  are integrable on  $[a, b]$  and  $k$  is a constant, then the functions of  $kf$  and  $f \pm g$  are integrable on  $[a, b]$ , and

1.  $\int_a^b kf(x) dx = k \int_a^b f(x) dx$
2.  $\int_a^b [f(x) \pm g(x)] dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx.$

Note that Property 2 of Theorem 4.7 can be extended to cover any finite number of functions. For example,

$$\int_a^b [f(x) + g(x) + h(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx + \int_a^b h(x) dx.$$

### EXAMPLE 6 Evaluation of a Definite Integral

Evaluate  $\int_1^3 (-x^2 + 4x - 3) dx$  using each of the following values.

$$\int_1^3 x^2 dx = \frac{26}{3}, \quad \int_1^3 x dx = 4, \quad \int_1^3 dx = 2$$

**Solution**

$$\begin{aligned} \int_1^3 (-x^2 + 4x - 3) dx &= \int_1^3 (-x^2) dx + \int_1^3 4x dx + \int_1^3 (-3) dx \\ &= -\int_1^3 x^2 dx + 4 \int_1^3 x dx - 3 \int_1^3 dx \\ &= -\left(\frac{26}{3}\right) + 4(4) - 3(2) \\ &= \frac{4}{3} \end{aligned}$$




Figure 4.26

If  $f$  and  $g$  are continuous on the closed interval  $[a, b]$  and

$$0 \leq f(x) \leq g(x)$$

for  $a \leq x \leq b$ , the following properties are true. First, the area of the region bounded by the graph of  $f$  and the  $x$ -axis (between  $a$  and  $b$ ) must be nonnegative. Second, this area must be less than or equal to the area of the region bounded by the graph of  $g$  and the  $x$ -axis (between  $a$  and  $b$ ), as shown in Figure 4.26. These two results are generalized in Theorem 4.8. (A proof of this theorem is given in Appendix A.)

**THEOREM 4.8 Preservation of Inequality**

1. If  $f$  is integrable and nonnegative on the closed interval  $[a, b]$ , then

$$0 \leq \int_a^b f(x) dx.$$

2. If  $f$  and  $g$  are integrable on the closed interval  $[a, b]$  and  $f(x) \leq g(x)$  for every  $x$  in  $[a, b]$ , then

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

**Exercises for Section 4.3**

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.

In Exercises 1 and 2, use Example 1 as a model to evaluate the limit

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(c_i) \Delta x_i$$

over the region bounded by the graphs of the equations.

1.  $f(x) = \sqrt{x}$ ,  $y = 0$ ,  $x = 0$ ,  $x = 3$

(Hint: Let  $c_i = 3i^2/n^2$ .)

2.  $f(x) = \sqrt[3]{x}$ ,  $y = 0$ ,  $x = 0$ ,  $x = 1$

(Hint: Let  $c_i = i^3/n^3$ .)

In Exercises 3–8, evaluate the definite integral by the limit definition.

3.  $\int_4^{10} 6 dx$

4.  $\int_{-2}^3 x dx$

5.  $\int_{-1}^1 x^3 dx$

6.  $\int_1^3 3x^2 dx$

7.  $\int_1^2 (x^2 + 1) dx$

8.  $\int_{-1}^2 (3x^2 + 2) dx$

In Exercises 9–12, write the limit as a definite integral on the interval  $[a, b]$ , where  $c_i$  is any point in the  $i$ th subinterval.

Limit

Interval

9.  $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n (3c_i + 10) \Delta x_i$

$[-1, 5]$

10.  $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n 6c_i(4 - c_i)^2 \Delta x_i$

$[0, 4]$

11.  $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n \sqrt{c_i^2 + 4} \Delta x_i$


$[0, 3]$

12.  $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n \left( \frac{3}{c_i^2} \right) \Delta x_i$


$[1, 3]$

In Exercises 13–22, set up a definite integral that yields the area of the region. (Do not evaluate the integral.)


13.  $f(x) = 3$




14.  $f(x) = 4 - 2x$




15.  $f(x) = 4 - |x|$




16.  $f(x) = x^2$




17.  $f(x) = 4 - x^2$




18.  $f(x) = \frac{1}{x^2 + 1}$




19.  $f(x) = \sin x$




20.  $f(x) = \tan x$



21.  $g(y) = y^3$



22.  $f(y) = (y - 2)^2$



In Exercises 23–32, sketch the region whose area is given by the definite integral. Then use a geometric formula to evaluate the integral ( $a > 0, r > 0$ ).

23.  $\int_0^3 4 \, dx$

24.  $\int_{-a}^a 4 \, dx$

25.  $\int_0^4 x \, dx$

26.  $\int_0^4 \frac{x}{2} \, dx$

27.  $\int_0^2 (2x + 5) \, dx$

28.  $\int_0^8 (8 - x) \, dx$

29.  $\int_{-1}^1 (1 - |x|) \, dx$

30.  $\int_{-a}^a (a - |x|) \, dx$

31.  $\int_{-3}^3 \sqrt{9 - x^2} \, dx$

32.  $\int_{-r}^r \sqrt{r^2 - x^2} \, dx$

In Exercises 33–40, evaluate the integral using the following values.

$$\int_2^4 x^3 \, dx = 60, \quad \int_2^4 x \, dx = 6, \quad \int_2^4 dx = 2$$

33.  $\int_4^2 x \, dx$

34.  $\int_2^2 x^3 \, dx$

35.  $\int_2^4 4x \, dx$

36.  $\int_2^4 15 \, dx$

37.  $\int_2^4 (x - 8) \, dx$

38.  $\int_2^4 (x^3 + 4) \, dx$

39.  $\int_2^4 \left(\frac{1}{2}x^3 - 3x + 2\right) \, dx$

40.  $\int_2^4 (6 + 2x - x^3) \, dx$

41. Given  $\int_0^5 f(x) \, dx = 10$  and  $\int_5^7 f(x) \, dx = 3$ , evaluate

(a)  $\int_0^7 f(x) \, dx$ .

(b)  $\int_5^0 f(x) \, dx$ .

(c)  $\int_5^5 f(x) \, dx$ .

(d)  $\int_0^5 3f(x) \, dx$ .

42. Given  $\int_0^3 f(x) \, dx = 4$  and  $\int_3^6 f(x) \, dx = -1$ , evaluate

(a)  $\int_0^6 f(x) \, dx$ .

(b)  $\int_6^3 f(x) \, dx$ .

(c)  $\int_3^3 f(x) \, dx$ .

(d)  $\int_3^6 -5f(x) \, dx$ .

43. Given  $\int_2^6 f(x) \, dx = 10$  and  $\int_2^6 g(x) \, dx = -2$ , evaluate

(a)  $\int_2^6 [f(x) + g(x)] \, dx$ .

(b)  $\int_2^6 [g(x) - f(x)] \, dx$ .

(c)  $\int_2^6 2g(x) \, dx$ .

(d)  $\int_2^6 3f(x) \, dx$ .

44. Given  $\int_{-1}^1 f(x) \, dx = 0$  and  $\int_0^1 f(x) \, dx = 5$ , evaluate

(a)  $\int_{-1}^0 f(x) \, dx$ .

(b)  $\int_0^1 f(x) \, dx - \int_{-1}^0 f(x) \, dx$ .

(c)  $\int_{-1}^1 3f(x) \, dx$ .

(d)  $\int_0^1 3f(x) \, dx$ .

45. Use the table of values to find lower and upper estimates of

$$\int_0^{10} f(x) \, dx.$$

Assume that  $f$  is a decreasing function.

$x$	0	2	4	6	8	10
$f(x)$	32	24	12	-4	-20	-36


46. Use the table of values to estimate

$$\int_0^6 f(x) \, dx.$$

Use three equal subintervals and the (a) left endpoints, (b) right endpoints, and (c) midpoints. If  $f$  is an increasing function, how does each estimate compare with the actual value? Explain your reasoning.


$x$	0	1	2	3	4	5	6
$f(x)$	-6	0	8	18	30	50	80

- 47. Think About It** The graph of  $f$  consists of line segments and a semicircle, as shown in the figure. Evaluate each definite integral by using geometric formulas.



- (a)  $\int_{-4}^2 f(x) dx$       (b)  $\int_2^6 f(x) dx$   
 (c)  $\int_{-4}^2 f(x) dx$       (d)  $\int_{-4}^6 f(x) dx$   
 (e)  $\int_{-4}^6 |f(x)| dx$       (f)  $\int_{-4}^6 [f(x) + 2] dx$

- 48. Think About It** The graph of  $f$  consists of line segments, as shown in the figure. Evaluate each definite integral by using geometric formulas.



- (a)  $\int_0^1 -f(x) dx$       (b)  $\int_3^4 3f(x) dx$   
 (c)  $\int_0^7 f(x) dx$       (d)  $\int_5^{11} f(x) dx$   
 (e)  $\int_0^{11} f(x) dx$       (f)  $\int_4^{10} f(x) dx$

- 49. Think About It** Consider the function  $f$  that is continuous on the interval  $[-5, 5]$  and for which

$$\int_0^5 f(x) dx = 4.$$

Evaluate each integral.


- (a)  $\int_0^5 [f(x) + 2] dx$   
 (b)  $\int_{-2}^3 f(x+2) dx$   
 (c)  $\int_{-5}^5 f(x) dx$  ( $f$  is even.)  
 (d)  $\int_{-5}^5 f(x) dx$  ( $f$  is odd.)

- 50. Think About It** A function  $f$  is defined below. Use geometric formulas to find  $\int_0^8 f(x) dx$ .

$$f(x) = \begin{cases} 4, & x < 4 \\ x, & x \geq 4 \end{cases}$$

### Writing About Concepts

In Exercises 51 and 52, use the figure to fill in the blank with the symbol  $<$ ,  $>$ , or  $=$ .



- 51.** The interval  $[1, 5]$  is partitioned into  $n$  subintervals of equal width  $\Delta x$ , and  $x_i$  is the left endpoint of the  $i$ th subinterval.

$$\sum_{i=1}^n f(x_i) \Delta x \quad \text{is} \quad \int_1^5 f(x) dx$$

- 52.** The interval  $[1, 5]$  is partitioned into  $n$  subintervals of equal width  $\Delta x$ , and  $x_i$  is the right endpoint of the  $i$ th subinterval.

$$\sum_{i=1}^n f(x_i) \Delta x \quad \text{is} \quad \int_1^5 f(x) dx$$

- 53.** Determine whether the function  $f(x) = \frac{1}{x-4}$  is integrable on the interval  $[3, 5]$ . Explain.

- 54.** Give an example of a function that is integrable on the interval  $[-1, 1]$ , but not continuous on  $[-1, 1]$ .

In Exercises 55–58, determine which value best approximates the definite integral. Make your selection on the basis of a sketch.

**55.**  $\int_0^4 \sqrt{x} dx$

- (a) 5      (b) -3      (c) 10      (d) 2      (e) 8

**56.**  $\int_0^{1/2} 4 \cos \pi x dx$

- (a) 4      (b)  $\frac{4}{3}$       (c) 16      (d)  $2\pi$       (e) -6

**57.**  $\int_0^1 2 \sin \pi x dx$

- (a) 6      (b)  $\frac{1}{2}$       (c) 4      (d)  $\frac{5}{4}$

**58.**  $\int_0^9 (1 + \sqrt{x}) dx$

- (a) -3      (b) 9      (c) 27      (d) 3

**Programming** Write a program for your graphing utility to approximate a definite integral using the Riemann sum

$$\sum_{i=1}^n f(c_i) \Delta x_i$$

where the subintervals are of equal width. The output should give three approximations of the integral where  $c_i$  is the left-hand endpoint  $L(n)$ , midpoint  $M(n)$ , and right-hand endpoint  $R(n)$  of each subinterval. In Exercises 59–62, use the program to approximate the definite integral and complete the table.

$n$	4	8	12	16	20
$L(n)$					
$M(n)$					
$R(n)$					

59.  $\int_0^3 x\sqrt{3-x} \, dx$

60.  $\int_0^3 \frac{5}{x^2+1} \, dx$

61.  $\int_0^{\pi/2} \sin^2 x \, dx$

62.  $\int_0^3 x \sin x \, dx$

**True or False?** In Exercises 63–68, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

63.  $\int_a^b [f(x) + g(x)] \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$

64.  $\int_a^b f(x)g(x) \, dx = \left[ \int_a^b f(x) \, dx \right] \left[ \int_a^b g(x) \, dx \right]$

65. If the norm of a partition approaches zero, then the number of subintervals approaches infinity.

66. If  $f$  is increasing on  $[a, b]$ , then the minimum value of  $f(x)$  on  $[a, b]$  is  $f(a)$ .

67. The value of

$$\int_a^b f(x) \, dx$$

must be positive.

68. The value of

$$\int_2^2 \sin(x^2) \, dx$$

is 0.

69. Find the Riemann sum for

$$f(x) = x^2 + 3x$$

over the interval  $[0, 8]$ , where  $x_0 = 0$ ,  $x_1 = 1$ ,  $x_2 = 3$ ,  $x_3 = 7$ , and  $x_4 = 8$ , and where  $c_1 = 1$ ,  $c_2 = 2$ ,  $c_3 = 5$ , and  $c_4 = 8$ .




Figure for 69




Figure for 70

70. Find the Riemann sum for  $f(x) = \sin x$  over the interval  $[0, 2\pi]$ , where  $x_0 = 0$ ,  $x_1 = \pi/4$ ,  $x_2 = \pi/3$ ,  $x_3 = \pi$ , and  $x_4 = 2\pi$ , and where  $c_1 = \pi/6$ ,  $c_2 = \pi/3$ ,  $c_3 = 2\pi/3$ , and  $c_4 = 3\pi/2$ .

71. Prove that  $\int_a^b x \, dx = \frac{b^2 - a^2}{2}$ .

72. Prove that  $\int_a^b x^2 \, dx = \frac{b^3 - a^3}{3}$ .


73. **Think About It** Determine whether the Dirichlet function

$$f(x) = \begin{cases} 1, & x \text{ is rational} \\ 0, & x \text{ is irrational} \end{cases}$$

is integrable on the interval  $[0, 1]$ . Explain.

74. Suppose the function  $f$  is defined on  $[0, 1]$ , as shown in the figure.

$$f(x) = \begin{cases} 0, & x = 0 \\ \frac{1}{x}, & 0 < x \leq 1 \end{cases}$$



Show that  $\int_0^1 f(x) \, dx$  does not exist. Why doesn't this contradict Theorem 4.4?

75. Find the constants  $a$  and  $b$  that maximize the value of

$$\int_a^b (1 - x^2) \, dx.$$

Explain your reasoning.

76. Evaluate, if possible, the integral  $\int_0^2 \lfloor x \rfloor \, dx$ .

77. Determine

$$\lim_{n \rightarrow \infty} \frac{1}{n^3} [1^2 + 2^2 + 3^2 + \cdots + n^2]$$

by using an appropriate Riemann sum.

## Section 4.4

## EXPLORATION

**Integration and Antidifferentiation**

Throughout this chapter, you have been using the integral sign to denote an antiderivative (a family of functions) and a definite integral (a number).

Antidifferentiation:  $\int f(x) dx$

Definite integration:  $\int_a^b f(x) dx$

The use of this same symbol for both operations makes it appear that they are related. In the early work with calculus, however, it was not known that the two operations were related. Do you think the symbol  $\int$  was first applied to antidifferentiation or to definite integration? Explain your reasoning. (*Hint:* The symbol was first used by Leibniz and was derived from the letter  $S$ .)


## The Fundamental Theorem of Calculus

- Evaluate a definite integral using the Fundamental Theorem of Calculus.
- Understand and use the Mean Value Theorem for Integrals.
- Find the average value of a function over a closed interval.
- Understand and use the Second Fundamental Theorem of Calculus.

## The Fundamental Theorem of Calculus

You have now been introduced to the two major branches of calculus: differential calculus (introduced with the tangent line problem) and integral calculus (introduced with the area problem). At this point, these two problems might seem unrelated—but there is a very close connection. The connection was discovered independently by Isaac Newton and Gottfried Leibniz and is stated in a theorem that is appropriately called the **Fundamental Theorem of Calculus**.

Informally, the theorem states that differentiation and (definite) integration are inverse operations, in the same sense that division and multiplication are inverse operations. To see how Newton and Leibniz might have anticipated this relationship, consider the approximations shown in Figure 4.27. The slope of the tangent line was defined using the *quotient*  $\Delta y/\Delta x$  (the slope of the secant line). Similarly, the area of a region under a curve was defined using the *product*  $\Delta y\Delta x$  (the area of a rectangle). So, at least in the primitive approximation stage, the operations of differentiation and definite integration appear to have an inverse relationship in the same sense that division and multiplication are inverse operations. The Fundamental Theorem of Calculus states that the limit processes (used to define the derivative and definite integral) preserve this inverse relationship.



Differentiation and definite integration have an “inverse” relationship.

Figure 4.27

**THEOREM 4.9 The Fundamental Theorem of Calculus**

If a function  $f$  is continuous on the closed interval  $[a, b]$  and  $F$  is an antiderivative of  $f$  on the interval  $[a, b]$ , then

$$\int_a^b f(x) dx = F(b) - F(a).$$



**Proof** The key to the proof is in writing the difference  $F(b) - F(a)$  in a convenient form. Let  $\Delta$  be the following partition of  $[a, b]$ .

$$a = x_0 < x_1 < x_2 < \cdots < x_{n-1} < x_n = b$$

By pairwise subtraction and addition of like terms, you can write

$$\begin{aligned} F(b) - F(a) &= F(x_n) - F(x_{n-1}) + F(x_{n-1}) - \cdots - F(x_1) + F(x_1) - F(x_0) \\ &= \sum_{i=1}^n [F(x_i) - F(x_{i-1})]. \end{aligned}$$

By the Mean Value Theorem, you know that there exists a number  $c_i$  in the  $i$ th subinterval such that

$$F'(c_i) = \frac{F(x_i) - F(x_{i-1})}{x_i - x_{i-1}}.$$

Because  $F'(c_i) = f(c_i)$ , you can let  $\Delta x_i = x_i - x_{i-1}$  and obtain

$$F(b) - F(a) = \sum_{i=1}^n f(c_i) \Delta x_i.$$

This important equation tells you that by applying the Mean Value Theorem you can always find a collection of  $c_i$ 's such that the *constant*  $F(b) - F(a)$  is a Riemann sum of  $f$  on  $[a, b]$ . Taking the limit (as  $\|\Delta\| \rightarrow 0$ ) produces

$$F(b) - F(a) = \int_a^b f(x) \, dx.$$

The following guidelines can help you understand the use of the Fundamental Theorem of Calculus.

### Guidelines for Using the Fundamental Theorem of Calculus

1. *Provided you can find* an antiderivative of  $f$ , you now have a way to evaluate a definite integral without having to use the limit of a sum.
2. When applying the Fundamental Theorem of Calculus, the following notation is convenient.

$$\begin{aligned} \int_a^b f(x) \, dx &= F(x) \Big|_a^b \\ &= F(b) - F(a) \end{aligned}$$

For instance, to evaluate  $\int_1^3 x^3 \, dx$ , you can write

$$\int_1^3 x^3 \, dx = \left[ \frac{x^4}{4} \right]_1^3 = \frac{3^4}{4} - \frac{1^4}{4} = \frac{81}{4} - \frac{1}{4} = 20.$$

3. It is not necessary to include a constant of integration  $C$  in the antiderivative because

$$\begin{aligned} \int_a^b f(x) \, dx &= \left[ F(x) + C \right]_a^b \\ &= [F(b) + C] - [F(a) + C] \\ &= F(b) - F(a). \end{aligned}$$

**EXAMPLE 1** Evaluating a Definite Integral

Evaluate each definite integral.


a.  $\int_1^2 (x^2 - 3) dx$       b.  $\int_1^4 3\sqrt{x} dx$       c.  $\int_0^{\pi/4} \sec^2 x dx$

**Solution**

a.  $\int_1^2 (x^2 - 3) dx = \left[ \frac{x^3}{3} - 3x \right]_1^2 = \left( \frac{8}{3} - 6 \right) - \left( \frac{1}{3} - 3 \right) = -\frac{2}{3}$

b.  $\int_1^4 3\sqrt{x} dx = 3 \int_1^4 x^{1/2} dx = 3 \left[ \frac{x^{3/2}}{3/2} \right]_1^4 = 2(4)^{3/2} - 2(1)^{3/2} = 14$

c.  $\int_0^{\pi/4} \sec^2 x dx = \tan x \Big|_0^{\pi/4} = 1 - 0 = 1$



The definite integral of  $y$  on  $[0, 2]$  is  $\frac{5}{2}$ .  
**Figure 4.28**

**EXAMPLE 2** A Definite Integral Involving Absolute ValueEvaluate  $\int_0^2 |2x - 1| dx$ .**Solution** Using Figure 4.28 and the definition of absolute value, you can rewrite the integrand as shown.

$$|2x - 1| = \begin{cases} -(2x - 1), & x < \frac{1}{2} \\ 2x - 1, & x \geq \frac{1}{2} \end{cases}$$

From this, you can rewrite the integral in two parts.

$$\begin{aligned} \int_0^2 |2x - 1| dx &= \int_0^{1/2} -(2x - 1) dx + \int_{1/2}^2 (2x - 1) dx \\ &= \left[ -x^2 + x \right]_0^{1/2} + \left[ x^2 - x \right]_{1/2}^2 \\ &= \left( -\frac{1}{4} + \frac{1}{2} \right) - (0 + 0) + (4 - 2) - \left( \frac{1}{4} - \frac{1}{2} \right) = \frac{5}{2} \end{aligned}$$

**EXAMPLE 3** Using the Fundamental Theorem to Find AreaFind the area of the region bounded by the graph of  $y = 2x^2 - 3x + 2$ , the  $x$ -axis, and the vertical lines  $x = 0$  and  $x = 2$ , as shown in Figure 4.29.**Solution** Note that  $y > 0$  on the interval  $[0, 2]$ .


$$\begin{aligned} \text{Area} &= \int_0^2 (2x^2 - 3x + 2) dx \\ &= \left[ \frac{2x^3}{3} - \frac{3x^2}{2} + 2x \right]_0^2 \\ &= \left( \frac{16}{3} - 6 + 4 \right) - (0 - 0 + 0) \\ &= \frac{10}{3} \end{aligned}$$

Integrate between  $x = 0$  and  $x = 2$ .


Find antiderivative.

Apply Fundamental Theorem.

Simplify.



The area of the region bounded by the graph of  $y$ , the  $x$ -axis,  $x = 0$ , and  $x = 2$  is  $\frac{10}{3}$ .  
**Figure 4.29**



Mean value rectangle:

$$f(c)(b-a) = \int_a^b f(x) dx$$

Figure 4.30

## The Mean Value Theorem for Integrals

In Section 4.2, you saw that the area of a region under a curve is greater than the area of an inscribed rectangle and less than the area of a circumscribed rectangle. The Mean Value Theorem for Integrals states that somewhere “between” the inscribed and circumscribed rectangles there is a rectangle whose area is precisely equal to the area of the region under the curve, as shown in Figure 4.30.

### THEOREM 4.10 Mean Value Theorem for Integrals

If  $f$  is continuous on the closed interval  $[a, b]$ , then there exists a number  $c$  in the closed interval  $[a, b]$  such that

$$\int_a^b f(x) dx = f(c)(b-a).$$

#### Proof

**Case 1:** If  $f$  is constant on the interval  $[a, b]$ , the theorem is clearly valid because  $c$  can be any point in  $[a, b]$ .

**Case 2:** If  $f$  is not constant on  $[a, b]$ , then, by the Extreme Value Theorem, you can choose  $f(m)$  and  $f(M)$  to be the minimum and maximum values of  $f$  on  $[a, b]$ . Because  $f(m) \leq f(x) \leq f(M)$  for all  $x$  in  $[a, b]$ , you can apply Theorem 4.8 to write the following.

$$\begin{aligned} \int_a^b f(m) dx &\leq \int_a^b f(x) dx \leq \int_a^b f(M) dx && \text{See Figure 4.31.} \\ f(m)(b-a) &\leq \int_a^b f(x) dx \leq f(M)(b-a) \\ f(m) &\leq \frac{1}{b-a} \int_a^b f(x) dx \leq f(M) \end{aligned}$$

From the third inequality, you can apply the Intermediate Value Theorem to conclude that there exists some  $c$  in  $[a, b]$  such that

$$f(c) = \frac{1}{b-a} \int_a^b f(x) dx \quad \text{or} \quad f(c)(b-a) = \int_a^b f(x) dx.$$





Figure 4.31

**NOTE** Notice that Theorem 4.10 does not specify how to determine  $c$ . It merely guarantees the existence of at least one number  $c$  in the interval.



$$\text{Average value} = \frac{1}{b-a} \int_a^b f(x) dx$$

Figure 4.32

## Average Value of a Function

The value of  $f(c)$  given in the Mean Value Theorem for Integrals is called the **average value** of  $f$  on the interval  $[a, b]$ .

### Definition of the Average Value of a Function on an Interval

If  $f$  is integrable on the closed interval  $[a, b]$ , then the **average value** of  $f$  on the interval is

$$\frac{1}{b-a} \int_a^b f(x) dx.$$

**NOTE** Notice in Figure 4.32 that the area of the region under the graph of  $f$  is equal to the area of the rectangle whose height is the average value.

To see why the average value of  $f$  is defined in this way, suppose that you partition  $[a, b]$  into  $n$  subintervals of equal width  $\Delta x = (b-a)/n$ . If  $c_i$  is any point in the  $i$ th subinterval, the arithmetic average (or mean) of the function values at the  $c_i$ 's is given by

$$a_n = \frac{1}{n} [f(c_1) + f(c_2) + \cdots + f(c_n)]. \quad \text{Average of } f(c_1), \dots, f(c_n)$$

By multiplying and dividing by  $(b-a)$ , you can write the average as

$$\begin{aligned} a_n &= \frac{1}{n} \sum_{i=1}^n f(c_i) \left( \frac{b-a}{b-a} \right) = \frac{1}{b-a} \sum_{i=1}^n f(c_i) \left( \frac{b-a}{n} \right) \\ &= \frac{1}{b-a} \sum_{i=1}^n f(c_i) \Delta x. \end{aligned}$$

Finally, taking the limit as  $n \rightarrow \infty$  produces the average value of  $f$  on the interval  $[a, b]$ , as given in the definition above.

This development of the average value of a function on an interval is only one of many practical uses of definite integrals to represent summation processes. In Chapter 7, you will study other applications, such as volume, arc length, centers of mass, and work.

### EXAMPLE 4 Finding the Average Value of a Function

Find the average value of  $f(x) = 3x^2 - 2x$  on the interval  $[1, 4]$ .

**Solution** The average value is given by

$$\begin{aligned} \frac{1}{b-a} \int_a^b f(x) dx &= \frac{1}{3} \int_1^4 (3x^2 - 2x) dx \\ &= \frac{1}{3} [x^3 - x^2]_1^4 \\ &= \frac{1}{3} [64 - 16 - (1 - 1)] = \frac{48}{3} = 16. \end{aligned}$$

(See Figure 4.33.)




Figure 4.33

George Hall/Corbis



The first person to fly at a speed greater than the speed of sound was Charles Yeager. On October 14, 1947, Yeager was clocked at 295.9 meters per second at an altitude of 12.2 kilometers. If Yeager had been flying at an altitude below 11.275 kilometers, this speed would not have “broken the sound barrier.” The photo above shows an F-14 *Tomcat*, a supersonic, twin-engine strike fighter. Currently, the *Tomcat* can reach heights of 15.24 kilometers and speeds up to 2 mach (707.78 meters per second).

### EXAMPLE 5 The Speed of Sound

At different altitudes in Earth’s atmosphere, sound travels at different speeds. The speed of sound  $s(x)$  (in meters per second) can be modeled by

$$s(x) = \begin{cases} -4x + 341, & 0 \leq x < 11.5 \\ 295, & 11.5 \leq x < 22 \\ \frac{3}{4}x + 278.5, & 22 \leq x < 32 \\ \frac{3}{2}x + 254.5, & 32 \leq x < 50 \\ -\frac{3}{2}x + 404.5, & 50 \leq x \leq 80 \end{cases}$$

where  $x$  is the altitude in kilometers (see Figure 4.34). What is the average speed of sound over the interval  $[0, 80]$ ?

**Solution** Begin by integrating  $s(x)$  over the interval  $[0, 80]$ . To do this, you can break the integral into five parts.


$$\begin{aligned} \int_0^{11.5} s(x) \, dx &= \int_0^{11.5} (-4x + 341) \, dx = \left[ -2x^2 + 341x \right]_0^{11.5} = 3657 \\ \int_{11.5}^{22} s(x) \, dx &= \int_{11.5}^{22} (295) \, dx = \left[ 295x \right]_{11.5}^{22} = 3097.5 \\ \int_{22}^{32} s(x) \, dx &= \int_{22}^{32} \left( \frac{3}{4}x + 278.5 \right) \, dx = \left[ \frac{3}{8}x^2 + 278.5x \right]_{22}^{32} = 2987.5 \\ \int_{32}^{50} s(x) \, dx &= \int_{32}^{50} \left( \frac{3}{2}x + 254.5 \right) \, dx = \left[ \frac{3}{4}x^2 + 254.5x \right]_{32}^{50} = 5688 \\ \int_{50}^{80} s(x) \, dx &= \int_{50}^{80} \left( -\frac{3}{2}x + 404.5 \right) \, dx = \left[ -\frac{3}{4}x^2 + 404.5x \right]_{50}^{80} = 9210 \end{aligned}$$

By adding the values of the five integrals, you have

$$\int_0^{80} s(x) \, dx = 24,640.$$

So, the average speed of sound from an altitude of 0 kilometers to an altitude of 80 kilometers is

$$\text{Average speed} = \frac{1}{80} \int_0^{80} s(x) \, dx = \frac{24,640}{80} = 308 \text{ meters per second.}$$




Speed of sound depends on altitude.

Figure 4.34


## The Second Fundamental Theorem of Calculus

Earlier you saw that the definite integral of  $f$  on the interval  $[a, b]$  was defined using the constant  $b$  as the upper limit of integration and  $x$  as the variable of integration. However, a slightly different situation may arise in which the variable  $x$  is used as the upper limit of integration. To avoid the confusion of using  $x$  in two different ways,  $t$  is temporarily used as the variable of integration. (Remember that the definite integral is *not* a function of its variable of integration.)

### The Definite Integral as a Number



### The Definite Integral as a Function of $x$



### EXPLORATION

Use a graphing utility to graph the function

$$F(x) = \int_0^x \cos t \, dt$$

for  $0 \leq x \leq \pi$ . Do you recognize this graph? Explain.

### EXAMPLE 6 The Definite Integral as a Function

Evaluate the function


$$F(x) = \int_0^x \cos t \, dt$$

at  $x = 0, \pi/6, \pi/4, \pi/3$ , and  $\pi/2$ .

**Solution** You could evaluate five different definite integrals, one for each of the given upper limits. However, it is much simpler to fix  $x$  (as a constant) temporarily to obtain

$$\int_0^x \cos t \, dt = \sin t \Big|_0^x = \sin x - \sin 0 = \sin x.$$

Now, using  $F(x) = \sin x$ , you can obtain the results shown in Figure 4.35.



$F(x) = \int_0^x \cos t \, dt$  is the area under the curve  $f(t) = \cos t$  from 0 to  $x$ .

Figure 4.35

You can think of the function  $F(x)$  as *accumulating* the area under the curve  $f(t) = \cos t$  from  $t = 0$  to  $t = x$ . For  $x = 0$ , the area is 0 and  $F(0) = 0$ . For  $x = \pi/2$ ,  $F(\pi/2) = 1$  gives the accumulated area under the cosine curve on the entire interval  $[0, \pi/2]$ . This interpretation of an integral as an **accumulation function** is used often in applications of integration.



In Example 6, note that the derivative of  $F$  is the original integrand (with only the variable changed). That is,

$$\frac{d}{dx}[F(x)] = \frac{d}{dx}[\sin x] = \frac{d}{dx}\left[\int_0^x \cos t \, dt\right] = \cos x.$$

This result is generalized in the following theorem, called the **Second Fundamental Theorem of Calculus**.

### THEOREM 4.11 The Second Fundamental Theorem of Calculus

If  $f$  is continuous on an open interval  $I$  containing  $a$ , then, for every  $x$  in the interval,

$$\frac{d}{dx}\left[\int_a^x f(t) \, dt\right] = f(x).$$

**Proof** Begin by defining  $F$  as

$$F(x) = \int_a^x f(t) \, dt.$$

Then, by the definition of the derivative, you can write

$$\begin{aligned} F'(x) &= \lim_{\Delta x \rightarrow 0} \frac{F(x + \Delta x) - F(x)}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\Delta x} \left[ \int_a^{x+\Delta x} f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\Delta x} \left[ \int_x^{x+\Delta x} f(t) \, dt + \int_a^x f(t) \, dt - \int_a^x f(t) \, dt \right] \\ &= \lim_{\Delta x \rightarrow 0} \frac{1}{\Delta x} \left[ \int_x^{x+\Delta x} f(t) \, dt \right]. \end{aligned}$$

From the Mean Value Theorem for Integrals (assuming  $\Delta x > 0$ ), you know there exists a number  $c$  in the interval  $[x, x + \Delta x]$  such that the integral in the expression above is equal to  $f(c) \Delta x$ . Moreover, because  $x \leq c \leq x + \Delta x$ , it follows that  $c \rightarrow x$  as  $\Delta x \rightarrow 0$ . So, you obtain

$$\begin{aligned} F'(x) &= \lim_{\Delta x \rightarrow 0} \left[ \frac{1}{\Delta x} f(c) \Delta x \right] \\ &= \lim_{\Delta x \rightarrow 0} f(c) \\ &= f(x). \end{aligned}$$

A similar argument can be made for  $\Delta x < 0$ .

**NOTE** Using the area model for definite integrals, you can view the approximation

$$f(x) \Delta x \approx \int_x^{x+\Delta x} f(t) \, dt$$

as saying that the area of the rectangle of height  $f(x)$  and width  $\Delta x$  is approximately equal to the area of the region lying between the graph of  $f$  and the  $x$ -axis on the interval  $[x, x + \Delta x]$ , as shown in Figure 4.36.




Figure 4.36

Note that the Second Fundamental Theorem of Calculus tells you that if a function is continuous, you can be sure that it has an antiderivative. This antiderivative need not, however, be an elementary function. (Recall the discussion of elementary functions in Section P.3.)

### EXAMPLE 7 Using the Second Fundamental Theorem of Calculus

Evaluate  $\frac{d}{dx} \left[ \int_0^x \sqrt{t^2 + 1} \, dt \right]$ .

**Solution** Note that  $f(t) = \sqrt{t^2 + 1}$  is continuous on the entire real line. So, using the Second Fundamental Theorem of Calculus, you can write

$$\frac{d}{dx} \left[ \int_0^x \sqrt{t^2 + 1} \, dt \right] = \sqrt{x^2 + 1}.$$

The differentiation shown in Example 7 is a straightforward application of the Second Fundamental Theorem of Calculus. The next example shows how this theorem can be combined with the Chain Rule to find the derivative of a function.

### EXAMPLE 8 Using the Second Fundamental Theorem of Calculus

Find the derivative of  $F(x) = \int_{\pi/2}^{x^3} \cos t \, dt$ .

**Solution** Using  $u = x^3$ , you can apply the Second Fundamental Theorem of Calculus with the Chain Rule as shown.

$$\begin{aligned} F'(x) &= \frac{dF}{du} \frac{du}{dx} && \text{Chain Rule} \\ &= \frac{d}{du} [F(x)] \frac{du}{dx} && \text{Definition of } \frac{dF}{du} \\ &= \frac{d}{du} \left[ \int_{\pi/2}^{x^3} \cos t \, dt \right] \frac{du}{dx} && \text{Substitute } \int_{\pi/2}^{x^3} \cos t \, dt \text{ for } F(x). \\ &= \frac{d}{du} \left[ \int_{\pi/2}^u \cos t \, dt \right] \frac{du}{dx} && \text{Substitute } u \text{ for } x^3. \\ &= (\cos u)(3x^2) && \text{Apply Second Fundamental Theorem of Calculus.} \\ &= (\cos x^3)(3x^2) && \text{Rewrite as function of } x. \end{aligned}$$

Because the integrand in Example 8 is easily integrated, you can verify the derivative as follows.

$$F(x) = \int_{\pi/2}^{x^3} \cos t \, dt = \sin t \Big|_{\pi/2}^{x^3} = \sin x^3 - \sin \frac{\pi}{2} = (\sin x^3) - 1$$

In this form, you can apply the Power Rule to verify that the derivative is the same as that obtained in Example 8.

$$F'(x) = (\cos x^3)(3x^2)$$

## Exercises for Section 4.4

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.

**Graphical Reasoning** In Exercises 1–4, use a graphing utility to graph the integrand. Use the graph to determine whether the definite integral is positive, negative, or zero.

1.  $\int_0^{\pi} \frac{4}{x^2 + 1} dx$

2.  $\int_0^{\pi} \cos x dx$

3.  $\int_{-2}^2 x\sqrt{x^2 + 1} dx$

4.  $\int_{-2}^2 x\sqrt{2-x} dx$

In Exercises 5–26, evaluate the definite integral of the algebraic function. Use a graphing utility to verify your result.

5.  $\int_0^1 2x dx$

6.  $\int_2^7 3 dv$

7.  $\int_{-1}^0 (x-2) dx$

8.  $\int_2^5 (-3v+4) dv$

9.  $\int_{-1}^1 (t^2-2) dt$

10.  $\int_1^3 (3x^2+5x-4) dx$

11.  $\int_0^1 (2t-1)^2 dt$

12.  $\int_{-1}^1 (t^3-9t) dt$

13.  $\int_1^2 \left(\frac{3}{x^2}-1\right) dx$

14.  $\int_{-2}^{-1} \left(u-\frac{1}{u^2}\right) du$

15.  $\int_1^4 \frac{u-2}{\sqrt{u}} du$

16.  $\int_{-3}^3 v^{1/3} dv$

17.  $\int_{-1}^1 (\sqrt[3]{t}-2) dt$

18.  $\int_1^8 \sqrt{\frac{2}{x}} dx$

19.  $\int_0^1 \frac{x-\sqrt{x}}{3} dx$

20.  $\int_0^2 (2-t)\sqrt{t} dt$

21.  $\int_{-1}^0 (t^{1/3}-t^{2/3}) dt$

22.  $\int_{-8}^{-1} \frac{x-x^2}{2\sqrt[3]{x}} dx$

23.  $\int_0^3 |2x-3| dx$

24.  $\int_1^4 (3-|x-3|) dx$

25.  $\int_0^3 |x^2-4| dx$

26.  $\int_0^4 |x^2-4x+3| dx$

In Exercises 27–32, evaluate the definite integral of the trigonometric function. Use a graphing utility to verify your result.

27.  $\int_0^{\pi} (1+\sin x) dx$

28.  $\int_0^{\pi/4} \frac{1-\sin^2 \theta}{\cos^2 \theta} d\theta$

29.  $\int_{-\pi/6}^{\pi/6} \sec^2 x dx$


30.  $\int_{\pi/4}^{\pi/2} (2-\csc^2 x) dx$

31.  $\int_{-\pi/3}^{\pi/3} 4 \sec \theta \tan \theta d\theta$


32.  $\int_{-\pi/2}^{\pi/2} (2t+\cos t) dt$

In Exercises 33–38, determine the area of the given region.


33.  $y = x - x^2$




34.  $y = 1 - x^4$




35.  $y = (3-x)\sqrt{x}$




36.  $y = \frac{1}{x^2}$



37.  $y = \cos x$



38.  $y = x + \sin x$



In Exercises 39–42, find the area of the region bounded by the graphs of the equations.

39.  $y = 3x^2 + 1$ ,  $x = 0$ ,  $x = 2$ ,  $y = 0$

40.  $y = 1 + \sqrt[3]{x}$ ,  $x = 0$ ,  $x = 8$ ,  $y = 0$

41.  $y = x^3 + x$ ,  $x = 2$ ,  $y = 0$

42.  $y = -x^2 + 3x$ ,  $y = 0$

In Exercises 43–46, find the value(s) of  $c$  guaranteed by the Mean Value Theorem for Integrals for the function over the given interval.

43.  $f(x) = x - 2\sqrt{x}$ ,  $[0, 2]$

44.  $f(x) = \frac{9}{x^3}$ ,  $[1, 3]$

45.  $f(x) = 2 \sec^2 x$ ,  $[-\pi/4, \pi/4]$

46.  $f(x) = \cos x$ ,  $[-\pi/3, \pi/3]$

In Exercises 47–50, find the average value of the function over the given interval and all values of  $x$  in the interval for which the function equals its average value.

47.  $f(x) = 4 - x^2$ ,  $[-2, 2]$

48.  $f(x) = \frac{4(x^2+1)}{x^2}$ ,  $[1, 3]$

49.  $f(x) = \sin x$ ,  $[0, \pi]$

50.  $f(x) = \cos x$ ,  $[0, \pi/2]$

- 51. Velocity** The graph shows the velocity, in feet per second, of a car accelerating from rest. Use the graph to estimate the distance the car travels in 8 seconds.




Figure for 51




Figure for 52

- 52. Velocity** The graph shows the velocity of a car as soon as the driver applies the brakes. Use the graph to estimate how far the car travels before it comes to a stop.

### Writing About Concepts

- 53.** State the Fundamental Theorem of Calculus.

- 54.** The graph of  $f$  is shown in the figure.

- Evaluate  $\int_1^7 f(x) dx$ .
- Determine the average value of  $f$  on the interval  $[1, 7]$ .
- Determine the answers to parts (a) and (b) if the graph is translated two units upward.




Figure for 54




Figure for 55–60

In Exercises 55–60, use the graph of  $f$  shown in the figure. The shaded region  $A$  has an area of 1.5, and  $\int_0^6 f(x) dx = 3.5$ . Use this information to fill in the blanks.

- $\int_0^2 f(x) dx =$
- $\int_2^6 f(x) dx =$
- $\int_0^6 |f(x)| dx =$
- $\int_0^2 -2f(x) dx =$
- $\int_0^6 [2 + f(x)] dx =$
- The average value of  $f$  over the interval  $[0, 6]$  is .

- 61. Force** The force  $F$  (in newtons) of a hydraulic cylinder in a press is proportional to the square of  $\sec x$ , where  $x$  is the distance (in meters) that the cylinder is extended in its cycle. The domain of  $F$  is  $[0, \pi/3]$ , and  $F(0) = 500$ .

- Find  $F$  as a function of  $x$ .
- Find the average force exerted by the press over the interval  $[0, \pi/3]$ .

- 62. Blood Flow** The velocity  $v$  of the flow of blood at a distance  $r$  from the central axis of an artery of radius  $R$  is

$$v = k(R^2 - r^2)$$

where  $k$  is the constant of proportionality. Find the average rate of flow of blood along a radius of the artery. (Use 0 and  $R$  as the limits of integration.)

- 63. Respiratory Cycle** The volume  $V$  in liters of air in the lungs during a five-second respiratory cycle is approximated by the model

$$V = 0.1729t + 0.1522t^2 - 0.0374t^3$$

where  $t$  is the time in seconds. Approximate the average volume of air in the lungs during one cycle.



- 64. Average Sales** A company fits a model to the monthly sales data of a seasonal product. The model is

$$S(t) = \frac{t}{4} + 1.8 + 0.5 \sin\left(\frac{\pi t}{6}\right), \quad 0 \leq t \leq 24$$

where  $S$  is sales (in thousands) and  $t$  is time in months.

- Use a graphing utility to graph  $f(t) = 0.5 \sin(\pi t/6)$  for  $0 \leq t \leq 24$ . Use the graph to explain why the average value of  $f(t)$  is 0 over the interval.
- Use a graphing utility to graph  $S(t)$  and the line  $g(t) = t/4 + 1.8$  in the same viewing window. Use the graph and the result of part (a) to explain why  $g$  is called the *trend line*.




- 65. Modeling Data** An experimental vehicle is tested on a straight track. It starts from rest, and its velocity  $v$  (meters per second) is recorded in the table every 10 seconds for 1 minute.

$t$	0	10	20	30	40	50	60
$v$	0	5	21	40	62	78	83

- Use a graphing utility to find a model of the form  $v = at^3 + bt^2 + ct + d$  for the data.
- Use a graphing utility to plot the data and graph the model.
- Use the Fundamental Theorem of Calculus to approximate the distance traveled by the vehicle during the test.



- (b) Plot the points from the table in part (a) and graph  $g$ .  
 (c) Where does  $g$  have its minimum? Explain.  
 (d) Where does  $g$  have a maximum? Explain.  
 (e) On what interval does  $g$  increase at the greatest rate? Explain.  
 (f) Identify the zeros of  $g$ .



95. **Cost** The total cost  $C$  (in dollars) of purchasing and maintaining a piece of equipment for  $x$  years is

$$C(x) = 5000 \left( 25 + 3 \int_0^x t^{1/4} dt \right).$$

- (a) Perform the integration to write  $C$  as a function of  $x$ .  
 (b) Find  $C(1)$ ,  $C(5)$ , and  $C(10)$ .  
 96. **Area** The area  $A$  between the graph of the function  $g(t) = 4 - 4/t^2$  and the  $t$ -axis over the interval  $[1, x]$  is

$$A(x) = \int_1^x \left( 4 - \frac{4}{t^2} \right) dt.$$

- (a) Find the horizontal asymptote of the graph of  $g$ .  
 (b) Integrate to find  $A$  as a function of  $x$ . Does the graph of  $A$  have a horizontal asymptote? Explain.

**Rectilinear Motion** In Exercises 97–99, consider a particle moving along the  $x$ -axis where  $x(t)$  is the position of the particle at time  $t$ ,  $x'(t)$  is its velocity, and  $\int_a^b |x'(t)| dt$  is the distance the particle travels in the interval of time.


97. The position function is given by  $x(t) = t^3 - 6t^2 + 9t - 2$ ,  $0 \leq t \leq 5$ . Find the total distance the particle travels in 5 units of time.  
 98. Repeat Exercise 97 for the position function given by  $x(t) = (t - 1)(t - 3)^2$ ,  $0 \leq t \leq 5$ .

99. A particle moves along the  $x$ -axis with velocity  $v(t) = 1/\sqrt{t}$ ,  $t > 0$ . At time  $t = 1$ , its position is  $x = 4$ . Find the total distance traveled by the particle on the interval  $1 \leq t \leq 4$ .

100. **Buffon's Needle Experiment** A horizontal plane is ruled with parallel lines 2 inches apart. A two-inch needle is tossed randomly onto the plane. The probability that the needle will touch a line is

$$P = \frac{2}{\pi} \int_0^{\pi/2} \sin \theta d\theta$$

where  $\theta$  is the acute angle between the needle and any one of the parallel lines. Find this probability.



**True or False?** In Exercises 101 and 102, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

101. If  $F'(x) = G'(x)$  on the interval  $[a, b]$ , then  $F(b) - F(a) = G(b) - G(a)$ .  
 102. If  $f$  is continuous on  $[a, b]$ , then  $f$  is integrable on  $[a, b]$ .  
 103. **Find the Error** Describe why the statement is incorrect.

$$\int_{-1}^1 x^{-2} dx = \left[ -x^{-1} \right]_{-1}^1 = (-1) - 1 = -2$$

104. Prove that  $\frac{d}{dx} \left[ \int_{u(x)}^{v(x)} f(t) dt \right] = f(v(x))v'(x) - f(u(x))u'(x)$ .  
 105. Show that the function

$$f(x) = \int_0^{1/x} \frac{1}{t^2 + 1} dt + \int_0^x \frac{1}{t^2 + 1} dt$$

is constant for  $x > 0$ .

106. Let  $G(x) = \int_0^x \left[ s \int_0^s f(t) dt \right] ds$ , where  $f$  is continuous for all real  $t$ . Find (a)  $G(0)$ , (b)  $G'(0)$ , (c)  $G''(x)$ , and (d)  $G''(0)$ .

## Section Project: Demonstrating the Fundamental Theorem

Use a graphing utility to graph the function  $y_1 = \sin^2 t$  on the interval  $0 \leq t \leq \pi$ . Let  $F(x)$  be the following function of  $x$ .

$$F(x) = \int_0^x \sin^2 t dt$$

- (a) Complete the table. Explain why the values of  $F$  are increasing.

$x$	0	$\pi/6$	$\pi/3$	$\pi/2$	$2\pi/3$	$5\pi/6$	$\pi$
$F(x)$							

- (b) Use the integration capabilities of a graphing utility to graph  $F$ .  
 (c) Use the differentiation capabilities of a graphing utility to graph  $F'(x)$ . How is this graph related to the graph in part (b)?  
 (d) Verify that the derivative of  $y = (1/2)t - (\sin 2t)/4$  is  $\sin^2 t$ . Graph  $y$  and write a short paragraph about how this graph is related to those in parts (b) and (c).



## Section 4.5

## Integration by Substitution

- Use pattern recognition to find an indefinite integral.
- Use a change of variables to find an indefinite integral.
- Use the General Power Rule for Integration to find an indefinite integral.
- Use a change of variables to evaluate a definite integral.
- Evaluate a definite integral involving an even or odd function.

## Pattern Recognition

In this section you will study techniques for integrating composite functions. The discussion is split into two parts—*pattern recognition* and *change of variables*. Both techniques involve a ***u*-substitution**. With pattern recognition you perform the substitution mentally, and with change of variables you write the substitution steps.

The role of substitution in integration is comparable to the role of the Chain Rule in differentiation. Recall that for differentiable functions given by  $y = F(u)$  and  $u = g(x)$ , the Chain Rule states that

$$\frac{d}{dx}[F(g(x))] = F'(g(x))g'(x).$$

From the definition of an antiderivative, it follows that

$$\begin{aligned}\int F'(g(x))g'(x) dx &= F(g(x)) + C \\ &= F(u) + C.\end{aligned}$$

These results are summarized in the following theorem.

**THEOREM 4.12** Antidifferentiation of a Composite Function

Let  $g$  be a function whose range is an interval  $I$ , and let  $f$  be a function that is continuous on  $I$ . If  $g$  is differentiable on its domain and  $F$  is an antiderivative of  $f$  on  $I$ , then

$$\int f(g(x))g'(x) dx = F(g(x)) + C.$$

If  $u = g(x)$ , then  $du = g'(x) dx$  and

$$\int f(u) du = F(u) + C.$$

**NOTE** The statement of Theorem 4.12 doesn't tell how to distinguish between  $f(g(x))$  and  $g'(x)$  in the integrand. As you become more experienced at integration, your skill in doing this will increase. Of course, part of the key is familiarity with derivatives.

**STUDY TIP** There are several techniques for applying substitution, each differing slightly from the others. However, you should remember that the goal is the same with every technique—you are trying to find an antiderivative of the integrand.

**EXPLORATION**


**Recognizing Patterns** The integrand in each of the following integrals fits the pattern  $f(g(x))g'(x)$ . Identify the pattern and use the result to evaluate the integral.

$$\text{a. } \int 2x(x^2 + 1)^4 dx \quad \text{b. } \int 3x^2 \sqrt{x^3 + 1} dx \quad \text{c. } \int \sec^2 x (\tan x + 3) dx$$

The next three integrals are similar to the first three. Show how you can multiply and divide by a constant to evaluate these integrals.

$$\text{d. } \int x(x^2 + 1)^4 dx \quad \text{e. } \int x^2 \sqrt{x^3 + 1} dx \quad \text{f. } \int 2 \sec^2 x (\tan x + 3) dx$$

Examples 1 and 2 show how to apply Theorem 4.12 *directly*, by recognizing the presence of  $f(g(x))$  and  $g'(x)$ . Note that the composite function in the integrand has an *outside function*  $f$  and an *inside function*  $g$ . Moreover, the derivative  $g'(x)$  is present as a factor of the integrand.



### EXAMPLE 1 Recognizing the $f(g(x))g'(x)$ Pattern

Find  $\int (x^2 + 1)^2(2x) dx$ .

**Solution** Letting  $g(x) = x^2 + 1$ , you obtain

$$g'(x) = 2x$$

and

$$f(g(x)) = f(x^2 + 1) = (x^2 + 1)^2.$$

From this, you can recognize that the integrand follows the  $f(g(x))g'(x)$  pattern. Using the Power Rule for Integration and Theorem 4.12, you can write

$$\int \overbrace{(x^2 + 1)^2}^{f(g(x))} \overbrace{(2x)}^{g'(x)} dx = \frac{1}{3} (x^2 + 1)^3 + C.$$

Try using the Chain Rule to check that the derivative of  $\frac{1}{3}(x^2 + 1)^3 + C$  is the integrand of the original integral.

### EXAMPLE 2 Recognizing the $f(g(x))g'(x)$ Pattern

Find  $\int 5 \cos 5x dx$ .

**Solution** Letting  $g(x) = 5x$ , you obtain

$$g'(x) = 5$$

and

$$f(g(x)) = f(5x) = \cos 5x.$$

From this, you can recognize that the integrand follows the  $f(g(x))g'(x)$  pattern. Using the Cosine Rule for Integration and Theorem 4.12, you can write

$$\int \overbrace{(\cos 5x)}^{f(g(x))} \overbrace{(5)}^{g'(x)} dx = \sin 5x + C.$$

You can check this by differentiating  $\sin 5x + C$  to obtain the original integrand.

**TECHNOLOGY** Try using a computer algebra system, such as *Maple*, *Derive*, *Mathematica*, *Mathcad*, or the *TI-89*, to solve the integrals given in Examples 1 and 2. Do you obtain the same antiderivatives that are listed in the examples?

The integrands in Examples 1 and 2 fit the  $f(g(x))g'(x)$  pattern exactly—you only had to recognize the pattern. You can extend this technique considerably with the Constant Multiple Rule

$$\int kf(x) \, dx = k \int f(x) \, dx.$$

Many integrands contain the essential part (the variable part) of  $g'(x)$  but are missing a constant multiple. In such cases, you can multiply and divide by the necessary constant multiple, as shown in Example 3.

### EXAMPLE 3 Multiplying and Dividing by a Constant

Find  $\int x(x^2 + 1)^2 \, dx$ .

**Solution** This is similar to the integral given in Example 1, except that the integrand is missing a factor of 2. Recognizing that  $2x$  is the derivative of  $x^2 + 1$ , you can let  $g(x) = x^2 + 1$  and supply the  $2x$  as follows.

$$\begin{aligned} \int x(x^2 + 1)^2 \, dx &= \int (x^2 + 1)^2 \left(\frac{1}{2}\right)(2x) \, dx && \text{Multiply and divide by 2.} \\ &= \frac{1}{2} \int \underbrace{(x^2 + 1)^2}_{f(g(x))} \underbrace{(2x)}_{g'(x)} \, dx && \text{Constant Multiple Rule} \\ &= \frac{1}{2} \left[ \frac{(x^2 + 1)^3}{3} \right] + C && \text{Integrate.} \\ &= \frac{1}{6} (x^2 + 1)^3 + C && \text{Simplify.} \end{aligned}$$

In practice, most people would not write as many steps as are shown in Example 3. For instance, you could evaluate the integral by simply writing

$$\begin{aligned} \int x(x^2 + 1)^2 \, dx &= \frac{1}{2} \int (x^2 + 1)^2 2x \, dx \\ &= \frac{1}{2} \left[ \frac{(x^2 + 1)^3}{3} \right] + C \\ &= \frac{1}{6} (x^2 + 1)^3 + C. \end{aligned}$$

**NOTE** Be sure you see that the *Constant Multiple Rule* applies only to *constants*. You cannot multiply and divide by a variable and then move the variable outside the integral sign. For instance,

$$\int (x^2 + 1)^2 \, dx \neq \frac{1}{2x} \int (x^2 + 1)^2 (2x) \, dx.$$

After all, if it were legitimate to move variable quantities outside the integral sign, you could move the entire integrand out and simplify the whole process. But the result would be incorrect.

## Change of Variables

With a formal **change of variables**, you completely rewrite the integral in terms of  $u$  and  $du$  (or any other convenient variable). Although this procedure can involve more written steps than the pattern recognition illustrated in Examples 1 to 3, it is useful for complicated integrands. The change of variable technique uses the Leibniz notation for the differential. That is, if  $u = g(x)$ , then  $du = g'(x) dx$ , and the integral in Theorem 4.12 takes the form

$$\int f(g(x))g'(x) dx = \int f(u) du = F(u) + C.$$

### EXAMPLE 4 Change of Variables

Find  $\int \sqrt{2x-1} dx$ .

**Solution** First, let  $u$  be the inner function,  $u = 2x - 1$ . Then calculate the differential  $du$  to be  $du = 2 dx$ . Now, using  $\sqrt{2x-1} = \sqrt{u}$  and  $dx = du/2$ , substitute to obtain

$$\begin{aligned} \int \sqrt{2x-1} dx &= \int \sqrt{u} \left( \frac{du}{2} \right) && \text{Integral in terms of } u \\ &= \frac{1}{2} \int u^{1/2} du && \text{Constant Multiple Rule} \\ &= \frac{1}{2} \left( \frac{u^{3/2}}{3/2} \right) + C && \text{Antiderivative in terms of } u \\ &= \frac{1}{3} u^{3/2} + C && \text{Simplify.} \\ &= \frac{1}{3} (2x-1)^{3/2} + C. && \text{Antiderivative in terms of } x \end{aligned}$$

**STUDY TIP** Because integration is usually more difficult than differentiation, you should always check your answer to an integration problem by differentiating. For instance, in Example 4 you should differentiate  $\frac{1}{3}(2x-1)^{3/2} + C$  to verify that you obtain the original integrand.



### EXAMPLE 5 Change of Variables

Find  $\int x\sqrt{2x-1} dx$ .

**Solution** As in the previous example, let  $u = 2x - 1$  and obtain  $dx = du/2$ . Because the integrand contains a factor of  $x$ , you must also solve for  $x$  in terms of  $u$ , as shown.

$$u = 2x - 1 \quad \Rightarrow \quad x = (u + 1)/2 \quad \text{Solve for } x \text{ in terms of } u.$$

Now, using substitution, you obtain

$$\begin{aligned} \int x\sqrt{2x-1} dx &= \int \left( \frac{u+1}{2} \right) u^{1/2} \left( \frac{du}{2} \right) \\ &= \frac{1}{4} \int (u^{3/2} + u^{1/2}) du \\ &= \frac{1}{4} \left( \frac{u^{5/2}}{5/2} + \frac{u^{3/2}}{3/2} \right) + C \\ &= \frac{1}{10} (2x-1)^{5/2} + \frac{1}{6} (2x-1)^{3/2} + C. \end{aligned}$$

To complete the change of variables in Example 5, you solved for  $x$  in terms of  $u$ . Sometimes this is very difficult. Fortunately it is not always necessary, as shown in the next example.

### EXAMPLE 6 Change of Variables

Find  $\int \sin^2 3x \cos 3x \, dx$ .

**Solution** Because  $\sin^2 3x = (\sin 3x)^2$ , you can let  $u = \sin 3x$ . Then

$$du = (\cos 3x)(3) \, dx.$$

Now, because  $\cos 3x \, dx$  is part of the original integral, you can write

$$\frac{du}{3} = \cos 3x \, dx.$$

Substituting  $u$  and  $du/3$  in the original integral yields

$$\begin{aligned} \int \sin^2 3x \cos 3x \, dx &= \int u^2 \frac{du}{3} \\ &= \frac{1}{3} \int u^2 \, du \\ &= \frac{1}{3} \left( \frac{u^3}{3} \right) + C \\ &= \frac{1}{9} \sin^3 3x + C. \end{aligned}$$

You can check this by differentiating.

$$\begin{aligned} \frac{d}{dx} \left[ \frac{1}{9} \sin^3 3x \right] &= \left( \frac{1}{9} \right) (3) (\sin 3x)^2 (\cos 3x) (3) \\ &= \sin^2 3x \cos 3x \end{aligned}$$

Because differentiation produces the original integrand, you know that you have obtained the correct antiderivative.

The steps used for integration by substitution are summarized in the following guidelines.

#### Guidelines for Making a Change of Variables

1. Choose a substitution  $u = g(x)$ . Usually, it is best to choose the *inner* part of a composite function, such as a quantity raised to a power.
2. Compute  $du = g'(x) \, dx$ .
3. Rewrite the integral in terms of the variable  $u$ .
4. Find the resulting integral in terms of  $u$ .
5. Replace  $u$  by  $g(x)$  to obtain an antiderivative in terms of  $x$ .
6. Check your answer by differentiating.

**STUDY TIP** When making a change of variables, be sure that your answer is written using the same variables as in the original integrand. For instance, in Example 6, you should not leave your answer as

$$\frac{1}{9} u^3 + C$$

but rather, replace  $u$  by  $\sin 3x$ .

## The General Power Rule for Integration

One of the most common  $u$ -substitutions involves quantities in the integrand that are raised to a power. Because of the importance of this type of substitution, it is given a special name—the **General Power Rule for Integration**. A proof of this rule follows directly from the (simple) Power Rule for Integration, together with Theorem 4.12.

### THEOREM 4.13 The General Power Rule for Integration

If  $g$  is a differentiable function of  $x$ , then

$$\int [g(x)]^n g'(x) dx = \frac{[g(x)]^{n+1}}{n+1} + C, \quad n \neq -1.$$

Equivalently, if  $u = g(x)$ , then

$$\int u^n du = \frac{u^{n+1}}{n+1} + C, \quad n \neq -1.$$

### EXAMPLE 7 Substitution and the General Power Rule

- a.  $\int 3(3x-1)^4 dx = \int \overbrace{(3x-1)^4}^{u^4} \overbrace{(3)}^{du} dx = \frac{\overbrace{(3x-1)^5}^{u^5/5}}{5} + C$
- b.  $\int (2x+1)(x^2+x) dx = \int \overbrace{(x^2+x)^1}^{u^1} \overbrace{(2x+1)}^{du} dx = \frac{\overbrace{(x^2+x)^2}^{u^2/2}}{2} + C$
- c.  $\int 3x^2 \sqrt{x^3-2} dx = \int \overbrace{(x^3-2)^{1/2}}^{u^{1/2}} \overbrace{(3x^2)}^{du} dx = \frac{\overbrace{(x^3-2)^{3/2}}^{u^{3/2}/(3/2)}}{3/2} + C = \frac{2}{3}(x^3-2)^{3/2} + C$
- d.  $\int \frac{-4x}{(1-2x^2)^2} dx = \int \overbrace{(1-2x^2)^{-2}}^{u^{-2}} \overbrace{(-4x)}^{du} dx = \frac{\overbrace{(1-2x^2)^{-1}}^{u^{-1}/(-1)}}{-1} + C = -\frac{1}{1-2x^2} + C$
- e.  $\int \cos^2 x \sin x dx = -\int \overbrace{(\cos x)^2}^{u^2} \overbrace{(-\sin x)}^{du} dx = -\frac{\overbrace{(\cos x)^3}^{u^3/3}}{3} + C$

#### EXPLORATION

Suppose you were asked to find one of the following integrals. Which one would you choose? Explain your reasoning.

a.  $\int \sqrt{x^3+1} dx$  or

$\int x^2 \sqrt{x^3+1} dx$

b.  $\int \tan(3x) \sec^2(3x) dx$  or

$\int \tan(3x) dx$

Some integrals whose integrands involve quantities raised to powers cannot be found by the General Power Rule. Consider the two integrals

$$\int x(x^2+1)^2 dx \quad \text{and} \quad \int (x^2+1)^2 dx.$$

The substitution  $u = x^2 + 1$  works in the first integral but not in the second. In the second, the substitution fails because the integrand lacks the factor  $x$  needed for  $du$ . Fortunately, for this particular integral, you can expand the integrand as  $(x^2+1)^2 = x^4 + 2x^2 + 1$  and use the (simple) Power Rule to integrate each term.



## Change of Variables for Definite Integrals

When using  $u$ -substitution with a definite integral, it is often convenient to determine the limits of integration for the variable  $u$  rather than to convert the antiderivative back to the variable  $x$  and evaluate at the original limits. This change of variables is stated explicitly in the next theorem. The proof follows from Theorem 4.12 combined with the Fundamental Theorem of Calculus.

### THEOREM 4.14 Change of Variables for Definite Integrals

If the function  $u = g(x)$  has a continuous derivative on the closed interval  $[a, b]$  and  $f$  is continuous on the range of  $g$ , then

$$\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du.$$

### EXAMPLE 8 Change of Variables

Evaluate  $\int_0^1 x(x^2 + 1)^3 dx$ .

**Solution** To evaluate this integral, let  $u = x^2 + 1$ . Then, you obtain

$$u = x^2 + 1 \Rightarrow du = 2x dx.$$

Before substituting, determine the new upper and lower limits of integration.

Lower Limit

$$\text{When } x = 0, u = 0^2 + 1 = 1.$$

Upper Limit

$$\text{When } x = 1, u = 1^2 + 1 = 2.$$

Now, you can substitute to obtain

$$\begin{aligned} \int_0^1 x(x^2 + 1)^3 dx &= \frac{1}{2} \int_0^1 (x^2 + 1)^3 (2x) dx && \text{Integration limits for } x \\ &= \frac{1}{2} \int_1^2 u^3 du && \text{Integration limits for } u \\ &= \frac{1}{2} \left[ \frac{u^4}{4} \right]_1^2 \\ &= \frac{1}{2} \left( 4 - \frac{1}{4} \right) \\ &= \frac{15}{8}. \end{aligned}$$

Try rewriting the antiderivative  $\frac{1}{2}(u^4/4)$  in terms of the variable  $x$  and evaluate the definite integral at the original limits of integration, as shown.

$$\begin{aligned} \frac{1}{2} \left[ \frac{u^4}{4} \right]_1^2 &= \frac{1}{2} \left[ \frac{(x^2 + 1)^4}{4} \right]_0^1 \\ &= \frac{1}{2} \left( 4 - \frac{1}{4} \right) = \frac{15}{8} \end{aligned}$$

Notice that you obtain the same result.

## Exercises for Section 4.5

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.

In Exercises 1–6, complete the table by identifying  $u$  and  $du$  for the integral.

$\int f(g(x))g'(x) dx$	$u = g(x)$	$du = g'(x) dx$
1. $\int (5x^2 + 1)^2(10x) dx$		
2. $\int x^2 \sqrt{x^3 + 1} dx$		
3. $\int \frac{x}{\sqrt{x^2 + 1}} dx$		
4. $\int \sec 2x \tan 2x dx$		
5. $\int \tan^2 x \sec^2 x dx$		
6. $\int \frac{\cos x}{\sin^2 x} dx$		

In Exercises 7–34, find the indefinite integral and check the result by differentiation.

7.  $\int (1 + 2x)^4(2) dx$
8.  $\int (x^2 - 9)^3(2x) dx$
9.  $\int \sqrt{9 - x^2}(-2x) dx$
10.  $\int \sqrt[3]{(1 - 2x^2)(-4x)} dx$
11.  $\int x^3(x^4 + 3)^2 dx$
12.  $\int x^2(x^3 + 5)^4 dx$
13.  $\int x^2(x^3 - 1)^4 dx$
14.  $\int x(4x^2 + 3)^3 dx$
15.  $\int t\sqrt{t^2 + 2} dt$
16.  $\int t^3\sqrt{t^4 + 5} dt$
17.  $\int 5x\sqrt[3]{1 - x^2} dx$
18.  $\int u^2\sqrt{u^3 + 2} du$
19.  $\int \frac{x}{(1 - x^2)^3} dx$
20.  $\int \frac{x^3}{(1 + x^4)^2} dx$
21.  $\int \frac{x^2}{(1 + x^3)^2} dx$
22.  $\int \frac{x^2}{(16 - x^3)^2} dx$
23.  $\int \frac{x}{\sqrt{1 - x^2}} dx$
24.  $\int \frac{x^3}{\sqrt{1 + x^4}} dx$
25.  $\int \left(1 + \frac{1}{t}\right)^3 \left(\frac{1}{t^2}\right) dt$
26.  $\int \left[x^2 + \frac{1}{(3x)^2}\right] dx$
27.  $\int \frac{1}{\sqrt{2x}} dx$
28.  $\int \frac{1}{2\sqrt{x}} dx$
29.  $\int \frac{x^2 + 3x + 7}{\sqrt{x}} dx$
30.  $\int \frac{t + 2t^2}{\sqrt{t}} dt$
31.  $\int t^2\left(t - \frac{2}{t}\right) dt$
32.  $\int \left(\frac{t^3}{3} + \frac{1}{4t^2}\right) dt$
33.  $\int (9 - y)\sqrt{y} dy$
34.  $\int 2\pi y(8 - y^{3/2}) dy$


In Exercises 35–38, solve the differential equation.

35.  $\frac{dy}{dx} = 4x + \frac{4x}{\sqrt{16 - x^2}}$
36.  $\frac{dy}{dx} = \frac{10x^2}{\sqrt{1 + x^3}}$
37.  $\frac{dy}{dx} = \frac{x + 1}{(x^2 + 2x - 3)^2}$
38.  $\frac{dy}{dx} = \frac{x - 4}{\sqrt{x^2 - 8x + 1}}$




**Slope Fields** In Exercises 39–42, a differential equation, a point, and a slope field are given. A *slope field* consists of line segments with slopes given by the differential equation. These line segments give a visual perspective of the directions of the solutions of the differential equation. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the given point. (To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).) (b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution. Compare the result with the sketches in part (a).


39.  $\frac{dy}{dx} = x\sqrt{4 - x^2}$   
(2, 2)




40.  $\frac{dy}{dx} = x^2(x^3 - 1)^2$   
(1, 0)



41.  $\frac{dy}{dx} = x \cos x^2$   
(0, 1)



42.  $\frac{dy}{dx} = -2 \sec(2x) \tan(2x)$   
(0, -1)



In Exercises 43–56, find the indefinite integral.

43.  $\int \pi \sin \pi x \, dx$

44.  $\int 4x^3 \sin x^4 \, dx$

45.  $\int \sin 2x \, dx$

46.  $\int \cos 6x \, dx$

47.  $\int \frac{1}{\theta^2} \cos \frac{1}{\theta} \, d\theta$

48.  $\int x \sin x^2 \, dx$

49.  $\int \sin 2x \cos 2x \, dx$

50.  $\int \sec(1-x) \tan(1-x) \, dx$

51.  $\int \tan^4 x \sec^2 x \, dx$

52.  $\int \sqrt{\tan x} \sec^2 x \, dx$

53.  $\int \frac{\csc^2 x}{\cot^3 x} \, dx$

54.  $\int \frac{\sin x}{\cos^3 x} \, dx$

55.  $\int \cot^2 x \, dx$

56.  $\int \csc^2\left(\frac{x}{2}\right) \, dx$

In Exercises 57–62, find an equation for the function  $f$  that has the given derivative and whose graph passes through the given point.

57.  $f'(x) = \cos \frac{x}{2}$

Point  $(0, 3)$

58.  $f'(x) = \pi \sec \pi x \tan \pi x$

Point  $\left(\frac{1}{3}, 1\right)$

59.  $f'(x) = \sin 4x$

Point  $\left(\frac{\pi}{4}, -\frac{3}{4}\right)$

60.  $f'(x) = \sec^2(2x)$

Point  $\left(\frac{\pi}{2}, 2\right)$

61.  $f'(x) = 2x(4x^2 - 10)^2$

Point  $(2, 10)$

62.  $f'(x) = -2x\sqrt{8-x^2}$

Point  $(2, 7)$

In Exercises 63–70, find the indefinite integral by the method shown in Example 5.

63.  $\int x\sqrt{x+2} \, dx$

64.  $\int x\sqrt{2x+1} \, dx$

65.  $\int x^2\sqrt{1-x} \, dx$

66.  $\int (x+1)\sqrt{2-x} \, dx$

67.  $\int \frac{x^2-1}{\sqrt{2x-1}} \, dx$

68.  $\int \frac{2x+1}{\sqrt{x+4}} \, dx$

69.  $\int \frac{-x}{(x+1)-\sqrt{x+1}} \, dx$

70.  $\int t\sqrt[3]{t-4} \, dt$

In Exercises 71–82, evaluate the definite integral. Use a graphing utility to verify your result.

71.  $\int_{-1}^1 x(x^2+1)^3 \, dx$

72.  $\int_{-2}^4 x^2(x^3+8)^2 \, dx$

73.  $\int_1^2 2x^2\sqrt{x^3+1} \, dx$

74.  $\int_0^1 x\sqrt{1-x^2} \, dx$

75.  $\int_0^4 \frac{1}{\sqrt{2x+1}} \, dx$

76.  $\int_0^2 \frac{x}{\sqrt{1+2x^2}} \, dx$

77.  $\int_1^9 \frac{1}{\sqrt{x}(1+\sqrt{x})^2} \, dx$

78.  $\int_0^2 x\sqrt[3]{4+x^2} \, dx$

79.  $\int_1^2 (x-1)\sqrt{2-x} \, dx$

80.  $\int_1^5 \frac{x}{\sqrt{2x-1}} \, dx$



81.  $\int_0^{\pi/2} \cos\left(\frac{2x}{3}\right) \, dx$

82.  $\int_{\pi/3}^{\pi/2} (x+\cos x) \, dx$

**Differential Equations** In Exercises 83–86, the graph of a function  $f$  is shown. Use the differential equation and the given point to find an equation of the function.



83.  $\frac{dy}{dx} = 18x^2(2x^3+1)^2$

84.  $\frac{dy}{dx} = \frac{-48}{(3x+5)^3}$



85.  $\frac{dy}{dx} = \frac{2x}{\sqrt{2x^2-1}}$



86.  $\frac{dy}{dx} = 4x + \frac{9x^2}{(3x^3+1)^{3/2}}$




In Exercises 87–92, find the area of the region. Use a graphing utility to verify your result.

87.  $\int_0^7 x\sqrt[3]{x+1} \, dx$


88.  $\int_{-2}^6 x^2\sqrt[3]{x+2} \, dx$




89.  $y = 2 \sin x + \sin 2x$




90.  $y = \sin x + \cos 2x$



91.  $\int_{\pi/2}^{2\pi/3} \sec^2\left(\frac{x}{2}\right) dx$



92.  $\int_{\pi/12}^{\pi/4} \csc 2x \cot 2x dx$



In Exercises 93–98, use a graphing utility to evaluate the integral. Graph the region whose area is given by the definite integral.

93.  $\int_0^4 \frac{x}{\sqrt{2x+1}} dx$

94.  $\int_0^2 x^3 \sqrt{x+2} dx$

95.  $\int_3^7 x \sqrt{x-3} dx$

96.  $\int_1^5 x^2 \sqrt{x-1} dx$

97.  $\int_0^3 \left( \theta + \cos \frac{\theta}{6} \right) d\theta$

98.  $\int_0^{\pi/2} \sin 2x dx$

**Writing** In Exercises 99 and 100, find the indefinite integral in two ways. Explain any difference in the forms of the answers.

99.  $\int (2x - 1)^2 dx$

100.  $\int \sin x \cos x dx$

In Exercises 101–104, evaluate the integral using the properties of even and odd functions as an aid.

101.  $\int_{-2}^2 x^2(x^2 + 1) dx$

102.  $\int_{-2}^2 x(x^2 + 1)^3 dx$

103.  $\int_{-\pi/2}^{\pi/2} \sin^2 x \cos x dx$

104.  $\int_{-\pi/2}^{\pi/2} \sin x \cos x dx$

105. Use  $\int_0^2 x^2 dx = \frac{8}{3}$  to evaluate each definite integral without using the Fundamental Theorem of Calculus.

(a)  $\int_{-2}^0 x^2 dx$

(b)  $\int_{-2}^2 x^2 dx$

(c)  $\int_0^2 -x^2 dx$

(d)  $\int_{-2}^0 3x^2 dx$

106. Use the symmetry of the graphs of the sine and cosine functions as an aid in evaluating each definite integral.

(a)  $\int_{-\pi/4}^{\pi/4} \sin x dx$

(b)  $\int_{-\pi/4}^{\pi/4} \cos x dx$

(c)  $\int_{-\pi/2}^{\pi/2} \cos x dx$

(d)  $\int_{-\pi/2}^{\pi/2} \sin x \cos x dx$

In Exercises 107 and 108, write the integral as the sum of the integral of an odd function and the integral of an even function. Use this simplification to evaluate the integral.

107.  $\int_{-4}^4 (x^3 + 6x^2 - 2x - 3) dx$

108.  $\int_{-\pi}^{\pi} (\sin 3x + \cos 3x) dx$

### Writing About Concepts

109. Describe why

$$\int x(5 - x^2)^3 dx \neq \int u^3 du$$

where  $u = 5 - x^2$ .

110. Without integrating, explain why

$$\int_{-2}^2 x(x^2 + 1)^2 dx = 0.$$

111. **Cash Flow** The rate of disbursement  $dQ/dt$  of a 2 million dollar federal grant is proportional to the square of  $100 - t$ . Time  $t$  is measured in days ( $0 \leq t \leq 100$ ), and  $Q$  is the amount that remains to be disbursed. Find the amount that remains to be disbursed after 50 days. Assume that all the money will be disbursed in 100 days.

112. **Depreciation** The rate of depreciation  $dV/dt$  of a machine is inversely proportional to the square of  $t + 1$ , where  $V$  is the value of the machine  $t$  years after it was purchased. The initial value of the machine was \$500,000, and its value decreased \$100,000 in the first year. Estimate its value after 4 years.

113. **Rainfall** The normal monthly rainfall at the Seattle-Tacoma airport can be approximated by the model

$$R = 3.121 + 2.399 \sin(0.524t + 1.377)$$

where  $R$  is measured in inches and  $t$  is the time in months, with  $t = 1$  corresponding to January. (Source: U.S. National Oceanic and Atmospheric Administration)

(a) Determine the extrema of the function over a one-year period.

(b) Use integration to approximate the normal annual rainfall. (Hint: Integrate over the interval  $[0, 12]$ .)

(c) Approximate the average monthly rainfall during the months of October, November, and December.

- 114. Sales** The sales  $S$  (in thousands of units) of a seasonal product are given by the model

$$S = 74.50 + 43.75 \sin \frac{\pi t}{6}$$

where  $t$  is the time in months, with  $t = 1$  corresponding to January. Find the average sales for each time period.

- The first quarter ( $0 \leq t \leq 3$ )
- The second quarter ( $3 \leq t \leq 6$ )
- The entire year ( $0 \leq t \leq 12$ )

- 115. Water Supply** A model for the flow rate of water at a pumping station on a given day is

$$R(t) = 53 + 7 \sin\left(\frac{\pi t}{6} + 3.6\right) + 9 \cos\left(\frac{\pi t}{12} + 8.9\right)$$

where  $0 \leq t \leq 24$ .  $R$  is the flow rate in thousands of gallons per hour, and  $t$  is the time in hours.



- Use a graphing utility to graph the rate function and approximate the maximum flow rate at the pumping station.
- Approximate the total volume of water pumped in 1 day.

- 116. Electricity** The oscillating current in an electrical circuit is

$$I = 2 \sin(60\pi t) + \cos(120\pi t)$$

where  $I$  is measured in amperes and  $t$  is measured in seconds. Find the average current for each time interval.

- $0 \leq t \leq \frac{1}{60}$
- $0 \leq t \leq \frac{1}{240}$
- $0 \leq t \leq \frac{1}{30}$

**Probability** In Exercises 117 and 118, the function

$$f(x) = kx^n(1-x)^m, \quad 0 \leq x \leq 1$$

where  $n > 0$ ,  $m > 0$ , and  $k$  is a constant, can be used to represent various probability distributions. If  $k$  is chosen such that

$$\int_0^1 f(x) dx = 1$$

the probability that  $x$  will fall between  $a$  and  $b$  ( $0 \leq a \leq b \leq 1$ ) is

$$P_{a,b} = \int_a^b f(x) dx.$$

- 117.** The probability that a person will remember between  $a\%$  and  $b\%$  of material learned in an experiment is

$$P_{a,b} = \int_a^b \frac{15}{4} x \sqrt{1-x} dx$$

where  $x$  represents the percent remembered. (See figure.)

- For a randomly chosen individual, what is the probability that he or she will recall between 50% and 75% of the material?

- What is the median percent recall? That is, for what value of  $b$  is it true that the probability of recalling 0 to  $b$  is 0.5?





Figure for 117

- 118.** The probability that ore samples taken from a region contain between  $a\%$  and  $b\%$  iron is

$$P_{a,b} = \int_a^b \frac{1155}{32} x^3(1-x)^{3/2} dx$$

where  $x$  represents the percent of iron. (See figure.) What is the probability that a sample will contain between

- 0% and 25% iron?
- 50% and 100% iron?




- 119. Temperature** The temperature in degrees Fahrenheit in a house is

$$T = 72 + 12 \sin\left[\frac{\pi(t-8)}{12}\right]$$

where  $t$  is time in hours, with  $t = 0$  representing midnight. The hourly cost of cooling a house is \$0.10 per degree.

- Find the cost  $C$  of cooling the house if its thermostat is set at 72°F by evaluating the integral


$$C = 0.1 \int_8^{20} \left[ 72 + 12 \sin \frac{\pi(t-8)}{12} - 72 \right] dt. \quad (\text{See figure.})$$



- (b) Find the savings from resetting the thermostat to 78°F by evaluating the integral

$$C = 0.1 \int_{10}^{18} \left[ 72 + 12 \sin \frac{\pi(t-8)}{12} - 78 \right] dt.$$

(See figure.)



- 120. Manufacturing** A manufacturer of fertilizer finds that national sales of fertilizer follow the seasonal pattern

$$F = 100,000 \left[ 1 + \sin \frac{2\pi(t-60)}{365} \right]$$

where  $F$  is measured in pounds and  $t$  represents the time in days, with  $t = 1$  corresponding to January 1. The manufacturer wants to set up a schedule to produce a uniform amount of fertilizer each day. What should this amount be?

- 121. Graphical Analysis** Consider the functions  $f$  and  $g$ , where

$$f(x) = 6 \sin x \cos^2 x \quad \text{and} \quad g(t) = \int_0^t f(x) dx.$$

- Use a graphing utility to graph  $f$  and  $g$  in the same viewing window.
- Explain why  $g$  is nonnegative.
- Identify the points on the graph of  $g$  that correspond to the extrema of  $f$ .
- Does each of the zeros of  $f$  correspond to an extremum of  $g$ ? Explain.
- Consider the function

$$h(t) = \int_{\pi/2}^t f(x) dx.$$

Use a graphing utility to graph  $h$ . What is the relationship between  $g$  and  $h$ ? Verify your conjecture.

- 122.** Find  $\lim_{n \rightarrow +\infty} \sum_{i=1}^n \frac{\sin(i\pi/n)}{n}$  by evaluating an appropriate definite integral over the interval  $[0, 1]$ .
- 123.** (a) Show that  $\int_0^1 x^2(1-x)^5 dx = \int_0^1 x^5(1-x)^2 dx$ .  
 (b) Show that  $\int_0^1 x^a(1-x)^b dx = \int_0^1 x^b(1-x)^a dx$ .
- 124.** (a) Show that  $\int_0^{\pi/2} \sin^2 x dx = \int_0^{\pi/2} \cos^2 x dx$ .  
 (b) Show that  $\int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n x dx$ , where  $n$  is a positive integer.

**True or False?** In Exercises 125–130, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

**125.**  $\int (2x+1)^2 dx = \frac{1}{3}(2x+1)^3 + C$

**126.**  $\int x(x^2+1) dx = \frac{1}{2}x^2(\frac{1}{3}x^3+x) + C$

**127.**  $\int_{-10}^{10} (ax^3 + bx^2 + cx + d) dx = 2 \int_0^{10} (bx^2 + d) dx$

**128.**  $\int_a^b \sin x dx = \int_a^{b+2\pi} \sin x dx$

**129.**  $4 \int \sin x \cos x dx = -\cos 2x + C$

**130.**  $\int \sin^2 2x \cos 2x dx = \frac{1}{3} \sin^3 2x + C$

- 131.** Assume that  $f$  is continuous everywhere and that  $c$  is a constant. Show that

$$\int_{ca}^{cb} f(x) dx = c \int_a^b f(cx) dx.$$

- 132.** (a) Verify that  $\sin u - u \cos u + C = \int u \sin u du$ .

(b) Use part (a) to show that  $\int_0^{\pi^2} \sin \sqrt{x} dx = 2\pi$ .

- 133.** Complete the proof of Theorem 4.15.

- 134.** Show that if  $f$  is continuous on the entire real number line, then

$$\int_a^b f(x+h) dx = \int_{a+h}^{b+h} f(x) dx.$$

### Putnam Exam Challenge

- 135.** If  $a_0, a_1, \dots, a_n$  are real numbers satisfying

$$\frac{a_0}{1} + \frac{a_1}{2} + \dots + \frac{a_n}{n+1} = 0$$

show that the equation  $a_0 + a_1x + a_2x^2 + \dots + a_nx^n = 0$  has at least one real zero.

- 136.** Find all the continuous positive functions  $f(x)$ , for  $0 \leq x \leq 1$ , such that

$$\int_0^1 f(x) dx = 1$$

$$\int_0^1 f(x)x dx = \alpha$$

$$\int_0^1 f(x)x^2 dx = \alpha^2$$

where  $\alpha$  is a real number.

These problems were composed by the Committee on the Putnam Prize Competition.  
 © The Mathematical Association of America. All rights reserved.



## Section 4.6

## Numerical Integration

- Approximate a definite integral using the Trapezoidal Rule.
- Approximate a definite integral using Simpson's Rule.
- Analyze the approximate errors in the Trapezoidal Rule and Simpson's Rule.

## The Trapezoidal Rule

Some elementary functions simply do not have antiderivatives that are elementary functions. For example, there is no elementary function that has any of the following functions as its derivative.

$$\sqrt[3]{x}\sqrt{1-x}, \quad \sqrt{x} \cos x, \quad \frac{\cos x}{x}, \quad \sqrt{1-x^3}, \quad \sin x^2$$

If you need to evaluate a definite integral involving a function whose antiderivative cannot be found, the Fundamental Theorem of Calculus cannot be applied, and you must resort to an approximation technique. Two such techniques are described in this section.

One way to approximate a definite integral is to use  $n$  trapezoids, as shown in Figure 4.41. In the development of this method, assume that  $f$  is continuous and positive on the interval  $[a, b]$ . So, the definite integral

$$\int_a^b f(x) \, dx$$

represents the area of the region bounded by the graph of  $f$  and the  $x$ -axis, from  $x = a$  to  $x = b$ . First, partition the interval  $[a, b]$  into  $n$  subintervals, each of width  $\Delta x = (b - a)/n$ , such that

$$a = x_0 < x_1 < x_2 < \cdots < x_n = b.$$

Then form a trapezoid for each subinterval (see Figure 4.42). The area of the  $i$ th trapezoid is

$$\text{Area of } i\text{th trapezoid} = \left[ \frac{f(x_{i-1}) + f(x_i)}{2} \right] \left( \frac{b - a}{n} \right).$$


This implies that the sum of the areas of the  $n$  trapezoids is

$$\begin{aligned} \text{Area} &= \left( \frac{b - a}{n} \right) \left[ \frac{f(x_0) + f(x_1)}{2} + \cdots + \frac{f(x_{n-1}) + f(x_n)}{2} \right] \\ &= \left( \frac{b - a}{2n} \right) [f(x_0) + f(x_1) + f(x_1) + f(x_2) + \cdots + f(x_{n-1}) + f(x_n)] \\ &= \left( \frac{b - a}{2n} \right) [f(x_0) + 2f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)]. \end{aligned}$$

Letting  $\Delta x = (b - a)/n$ , you can take the limit as  $n \rightarrow \infty$  to obtain


$$\begin{aligned} &\lim_{n \rightarrow \infty} \left( \frac{b - a}{2n} \right) [f(x_0) + 2f(x_1) + \cdots + 2f(x_{n-1}) + f(x_n)] \\ &= \lim_{n \rightarrow \infty} \left[ \frac{[f(a) - f(b)] \Delta x}{2} + \sum_{i=1}^n f(x_i) \Delta x \right] \\ &= \lim_{n \rightarrow \infty} \frac{[f(a) - f(b)](b - a)}{2n} + \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x \\ &= 0 + \int_a^b f(x) \, dx. \end{aligned}$$

The result is summarized in the following theorem.



The area of the region can be approximated using four trapezoids.

Figure 4.41



The area of the first trapezoid is  $\left[ \frac{f(x_0) + f(x_1)}{2} \right] \left( \frac{b - a}{n} \right)$ .

Figure 4.42

**THEOREM 4.16 The Trapezoidal Rule**

Let  $f$  be continuous on  $[a, b]$ . The Trapezoidal Rule for approximating  $\int_a^b f(x) dx$  is given by

$$\int_a^b f(x) dx \approx \frac{b-a}{2n} [f(x_0) + 2f(x_1) + 2f(x_2) + \cdots + 2f(x_{n-1}) + f(x_n)].$$

Moreover, as  $n \rightarrow \infty$ , the right-hand side approaches  $\int_a^b f(x) dx$ .

**NOTE** Observe that the coefficients in the Trapezoidal Rule have the following pattern.

$$1 \quad 2 \quad 2 \quad 2 \quad \cdots \quad 2 \quad 2 \quad 1$$

**EXAMPLE 1 Approximation with the Trapezoidal Rule**

Use the Trapezoidal Rule to approximate

$$\int_0^\pi \sin x dx.$$

Compare the results for  $n = 4$  and  $n = 8$ , as shown in Figure 4.43.


**Solution** When  $n = 4$ ,  $\Delta x = \pi/4$ , and you obtain

$$\begin{aligned} \int_0^\pi \sin x dx &\approx \frac{\pi}{8} \left( \sin 0 + 2 \sin \frac{\pi}{4} + 2 \sin \frac{\pi}{2} + 2 \sin \frac{3\pi}{4} + \sin \pi \right) \\ &= \frac{\pi}{8} (0 + \sqrt{2} + 2 + \sqrt{2} + 0) = \frac{\pi(1 + \sqrt{2})}{4} \approx 1.896. \end{aligned}$$


When  $n = 8$ ,  $\Delta x = \pi/8$ , and you obtain

$$\begin{aligned} \int_0^\pi \sin x dx &\approx \frac{\pi}{16} \left( \sin 0 + 2 \sin \frac{\pi}{8} + 2 \sin \frac{\pi}{4} + 2 \sin \frac{3\pi}{8} + 2 \sin \frac{\pi}{2} \right. \\ &\quad \left. + 2 \sin \frac{5\pi}{8} + 2 \sin \frac{3\pi}{4} + 2 \sin \frac{7\pi}{8} + \sin \pi \right) \\ &= \frac{\pi}{16} \left( 2 + 2\sqrt{2} + 4 \sin \frac{\pi}{8} + 4 \sin \frac{3\pi}{8} \right) \approx 1.974. \end{aligned}$$

For this particular integral, you could have found an antiderivative and determined that the exact area of the region is 2.



Four subintervals



Eight subintervals

Trapezoidal approximations  
Figure 4.43

**TECHNOLOGY** Most graphing utilities and computer algebra systems have built-in programs that can be used to approximate the value of a definite integral. Try using such a program to approximate the integral in Example 1. How close is your approximation?

When you use such a program, you need to be aware of its limitations. Often, you are given no indication of the degree of accuracy of the approximation. Other times, you may be given an approximation that is completely wrong. For instance, try using a built-in numerical integration program to evaluate

$$\int_{-1}^2 \frac{1}{x} dx.$$

Your calculator should give an error message. Does yours?

It is interesting to compare the Trapezoidal Rule with the Midpoint Rule given in Section 4.2 (Exercises 63–66). For the Trapezoidal Rule, you average the function values at the endpoints of the subintervals, but for the Midpoint Rule you take the function values of the subinterval midpoints.

$$\int_a^b f(x) dx \approx \sum_{i=1}^n f\left(\frac{x_i + x_{i-1}}{2}\right) \Delta x \quad \text{Midpoint Rule}$$

$$\int_a^b f(x) dx \approx \sum_{i=1}^n \left( \frac{f(x_i) + f(x_{i-1})}{2} \right) \Delta x \quad \text{Trapezoidal Rule}$$

**NOTE** There are two important points that should be made concerning the Trapezoidal Rule (or the Midpoint Rule). First, the approximation tends to become more accurate as  $n$  increases. For instance, in Example 1, if  $n = 16$ , the Trapezoidal Rule yields an approximation of 1.994. Second, although you could have used the Fundamental Theorem to evaluate the integral in Example 1, this theorem cannot be used to evaluate an integral as simple as  $\int_0^\pi \sin x^2 dx$  because  $\sin x^2$  has no elementary antiderivative. Yet, the Trapezoidal Rule can be applied easily to this integral.

### Simpson's Rule

One way to view the trapezoidal approximation of a definite integral is to say that on each subinterval you approximate  $f$  by a *first-degree* polynomial. In Simpson's Rule, named after the English mathematician Thomas Simpson (1710–1761), you take this procedure one step further and approximate  $f$  by *second-degree* polynomials.

Before presenting Simpson's Rule, we list a theorem for evaluating integrals of polynomials of degree 2 (or less).

#### THEOREM 4.17 Integral of $p(x) = Ax^2 + Bx + C$

If  $p(x) = Ax^2 + Bx + C$ , then

$$\int_a^b p(x) dx = \left( \frac{b-a}{6} \right) \left[ p(a) + 4p\left(\frac{a+b}{2}\right) + p(b) \right].$$

#### Proof


$$\begin{aligned} \int_a^b p(x) dx &= \int_a^b (Ax^2 + Bx + C) dx \\ &= \left[ \frac{Ax^3}{3} + \frac{Bx^2}{2} + Cx \right]_a^b \\ &= \frac{A(b^3 - a^3)}{3} + \frac{B(b^2 - a^2)}{2} + C(b - a) \\ &= \left( \frac{b-a}{6} \right) [2A(a^2 + ab + b^2) + 3B(b+a) + 6C] \end{aligned}$$

By expansion and collection of terms, the expression inside the brackets becomes

$$\underbrace{(Aa^2 + Ba + C)}_{p(a)} + 4 \underbrace{\left[ A\left(\frac{b+a}{2}\right)^2 + B\left(\frac{b+a}{2}\right) + C \right]}_{4p\left(\frac{a+b}{2}\right)} + \underbrace{(Ab^2 + Bb + C)}_{p(b)}$$

and you can write

$$\int_a^b p(x) dx = \left( \frac{b-a}{6} \right) \left[ p(a) + 4p\left(\frac{a+b}{2}\right) + p(b) \right].$$



$$\int_{x_0}^{x_2} p(x) dx \approx \int_{x_0}^{x_2} f(x) dx$$

Figure 4.44

To develop Simpson's Rule for approximating a definite integral, you again partition the interval  $[a, b]$  into  $n$  subintervals, each of width  $\Delta x = (b - a)/n$ . This time, however,  $n$  is required to be even, and the subintervals are grouped in pairs such that

$$a = x_0 < x_1 < x_2 < x_3 < x_4 < \cdots < x_{n-2} < x_{n-1} < x_n = b.$$

$[x_0, x_2]$ 
 $[x_2, x_4]$ 
 $[x_{n-2}, x_n]$

On each (double) subinterval  $[x_{i-2}, x_i]$ , you can approximate  $f$  by a polynomial  $p$  of degree less than or equal to 2. (See Exercise 55.) For example, on the subinterval  $[x_0, x_2]$ , choose the polynomial of least degree passing through the points  $(x_0, y_0)$ ,  $(x_1, y_1)$ , and  $(x_2, y_2)$ , as shown in Figure 4.44. Now, using  $p$  as an approximation of  $f$  on this subinterval, you have, by Theorem 4.17,

$$\begin{aligned} \int_{x_0}^{x_2} f(x) dx &\approx \int_{x_0}^{x_2} p(x) dx = \frac{x_2 - x_0}{6} \left[ p(x_0) + 4p\left(\frac{x_0 + x_2}{2}\right) + p(x_2) \right] \\ &= \frac{2[(b - a)/n]}{6} [p(x_0) + 4p(x_1) + p(x_2)] \\ &= \frac{b - a}{3n} [f(x_0) + 4f(x_1) + f(x_2)]. \end{aligned}$$

Repeating this procedure on the entire interval  $[a, b]$  produces the following theorem.

#### THEOREM 4.18 Simpson's Rule ( $n$ is even)

Let  $f$  be continuous on  $[a, b]$ . Simpson's Rule for approximating  $\int_a^b f(x) dx$  is

$$\begin{aligned} \int_a^b f(x) dx &\approx \frac{b - a}{3n} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \cdots \\ &\quad + 4f(x_{n-1}) + f(x_n)]. \end{aligned}$$

Moreover, as  $n \rightarrow \infty$ , the right-hand side approaches  $\int_a^b f(x) dx$ .

**NOTE** Observe that the coefficients in Simpson's Rule have the following pattern.

$$1 \quad 4 \quad 2 \quad 4 \quad 2 \quad 4 \quad \cdots \quad 4 \quad 2 \quad 4 \quad 1$$

In Example 1, the Trapezoidal Rule was used to estimate  $\int_0^\pi \sin x dx$ . In the next example, Simpson's Rule is applied to the same integral.



#### EXAMPLE 2 Approximation with Simpson's Rule

Use Simpson's Rule to approximate

$$\int_0^\pi \sin x dx.$$

Compare the results for  $n = 4$  and  $n = 8$ .

**Solution** When  $n = 4$ , you have

$$\int_0^\pi \sin x dx \approx \frac{\pi}{12} \left( \sin 0 + 4 \sin \frac{\pi}{4} + 2 \sin \frac{\pi}{2} + 4 \sin \frac{3\pi}{4} + \sin \pi \right) \approx 2.005.$$

When  $n = 8$ , you have  $\int_0^\pi \sin x dx \approx 2.0003$ .

**NOTE** In Example 1, the Trapezoidal Rule with  $n = 8$  approximated  $\int_0^\pi \sin x dx$  as 1.974. In Example 2, Simpson's Rule with  $n = 8$  gave an approximation of 2.0003. The antiderivative would produce the true value of 2.

## Error Analysis

If you must use an approximation technique, it is important to know how accurate you can expect the approximation to be. The following theorem, which is listed without proof, gives the formulas for estimating the errors involved in the use of Simpson's Rule and the Trapezoidal Rule.

### THEOREM 4.19 Errors in the Trapezoidal Rule and Simpson's Rule

If  $f$  has a continuous second derivative on  $[a, b]$ , then the error  $E$  in approximating  $\int_a^b f(x) dx$  by the Trapezoidal Rule is

$$E \leq \frac{(b-a)^3}{12n^2} [\max |f''(x)|], \quad a \leq x \leq b. \quad \text{Trapezoidal Rule}$$


Moreover, if  $f$  has a continuous fourth derivative on  $[a, b]$ , then the error  $E$  in approximating  $\int_a^b f(x) dx$  by Simpson's Rule is

$$E \leq \frac{(b-a)^5}{180n^4} [\max |f^{(4)}(x)|], \quad a \leq x \leq b. \quad \text{Simpson's Rule}$$

**TECHNOLOGY** If you have access to a computer algebra system, use it to evaluate the definite integral in Example 3. You should obtain a value of

$$\int_0^1 \sqrt{1+x^2} dx = \frac{1}{2} [\sqrt{2} + \ln(1+\sqrt{2})] \approx 1.14779.$$

("ln" represents the natural logarithmic function, which you will study in Section 5.1.)



$$1.144 \leq \int_0^1 \sqrt{1+x^2} dx \leq 1.164$$

Figure 4.45

Theorem 4.19 states that the errors generated by the Trapezoidal Rule and Simpson's Rule have upper bounds dependent on the extreme values of  $f''(x)$  and  $f^{(4)}(x)$  in the interval  $[a, b]$ . Furthermore, these errors can be made arbitrarily small by increasing  $n$ , provided that  $f''$  and  $f^{(4)}$  are continuous and therefore bounded in  $[a, b]$ .

### EXAMPLE 3 The Approximate Error in the Trapezoidal Rule

Determine a value of  $n$  such that the Trapezoidal Rule will approximate the value of  $\int_0^1 \sqrt{1+x^2} dx$  with an error that is less than 0.01.

**Solution** Begin by letting  $f(x) = \sqrt{1+x^2}$  and finding the second derivative of  $f$ .

$$f'(x) = x(1+x^2)^{-1/2} \quad \text{and} \quad f''(x) = (1+x^2)^{-3/2}$$

The maximum value of  $|f''(x)|$  on the interval  $[0, 1]$  is  $|f''(0)| = 1$ . So, by Theorem 4.19, you can write

$$E \leq \frac{(b-a)^3}{12n^2} |f''(0)| = \frac{1}{12n^2} (1) = \frac{1}{12n^2}.$$

To obtain an error  $E$  that is less than 0.01, you must choose  $n$  such that  $1/(12n^2) \leq 1/100$ .

$$100 \leq 12n^2 \quad \Rightarrow \quad n \geq \sqrt{\frac{100}{12}} \approx 2.89$$

So, you can choose  $n = 3$  (because  $n$  must be greater than or equal to 2.89) and apply the Trapezoidal Rule, as shown in Figure 4.45, to obtain

$$\begin{aligned} \int_0^1 \sqrt{1+x^2} dx &\approx \frac{1}{6} \left[ \sqrt{1+0^2} + 2\sqrt{1+\left(\frac{1}{3}\right)^2} + 2\sqrt{1+\left(\frac{2}{3}\right)^2} + \sqrt{1+1^2} \right] \\ &\approx 1.154. \end{aligned}$$

So, with an error no larger than 0.01, you know that

$$1.144 \leq \int_0^1 \sqrt{1+x^2} dx \leq 1.164.$$




## Exercises for Section 4.6

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.

In Exercises 1–10, use the Trapezoidal Rule and Simpson's Rule to approximate the value of the definite integral for the given value of  $n$ . Round your answer to four decimal places and compare the results with the exact value of the definite integral.

1.  $\int_0^2 x^2 dx$ ,  $n = 4$
2.  $\int_0^1 \left( \frac{x^2}{2} + 1 \right) dx$ ,  $n = 4$
3.  $\int_0^2 x^3 dx$ ,  $n = 4$
4.  $\int_1^2 \frac{1}{x^2} dx$ ,  $n = 4$
5.  $\int_0^2 x^3 dx$ ,  $n = 8$
6.  $\int_0^8 \sqrt[3]{x} dx$ ,  $n = 8$
7.  $\int_4^9 \sqrt{x} dx$ ,  $n = 8$
8.  $\int_1^3 (4 - x^2) dx$ ,  $n = 4$
9.  $\int_1^2 \frac{1}{(x+1)^2} dx$ ,  $n = 4$
10.  $\int_0^2 x\sqrt{x^2+1} dx$ ,  $n = 4$

 In Exercises 11–20, approximate the definite integral using the Trapezoidal Rule and Simpson's Rule with  $n = 4$ . Compare these results with the approximation of the integral using a graphing utility.

11.  $\int_0^2 \sqrt{1+x^3} dx$
12.  $\int_0^2 \frac{1}{\sqrt{1+x^3}} dx$
13.  $\int_0^1 \sqrt{x} \sqrt{1-x} dx$
14.  $\int_{\pi/2}^{\pi} \sqrt{x} \sin x dx$
15.  $\int_0^{\sqrt{\pi/2}} \cos x^2 dx$
16.  $\int_0^{\sqrt{\pi/4}} \tan x^2 dx$
17.  $\int_1^{1.1} \sin x^2 dx$
18.  $\int_0^{\pi/2} \sqrt{1+\cos^2 x} dx$
19.  $\int_0^{\pi/4} x \tan x dx$
20.  $\int_0^{\pi} f(x) dx$ ,  $f(x) = \begin{cases} \frac{\sin x}{x}, & x > 0 \\ 1, & x = 0 \end{cases}$

## Writing About Concepts

21. If the function  $f$  is concave upward on the interval  $[a, b]$ , will the Trapezoidal Rule yield a result greater than or less than  $\int_a^b f(x) dx$ ? Explain.
22. The Trapezoidal Rule and Simpson's Rule yield approximations of a definite integral  $\int_a^b f(x) dx$  based on polynomial approximations of  $f$ . What degree polynomial is used for each?


In Exercises 23–28, use the error formulas in Theorem 4.19 to estimate the error in approximating the integral, with  $n = 4$ , using (a) the Trapezoidal Rule and (b) Simpson's Rule.

23.  $\int_0^2 x^3 dx$
24.  $\int_1^3 (2x + 3) dx$

25.  $\int_0^1 \frac{1}{x+1} dx$
26.  $\int_2^4 \frac{1}{(x-1)^2} dx$
27.  $\int_0^{\pi} \cos x dx$
28.  $\int_0^1 \sin(\pi x) dx$

In Exercises 29–34, use the error formulas in Theorem 4.19 to find  $n$  such that the error in the approximation of the definite integral is less than 0.00001 using (a) the Trapezoidal Rule and (b) Simpson's Rule.

29.  $\int_1^3 \frac{1}{x} dx$
30.  $\int_0^1 \frac{1}{1+x} dx$
31.  $\int_0^2 \sqrt{x+2} dx$
32.  $\int_1^3 \frac{1}{\sqrt{x}} dx$
33.  $\int_0^1 \cos(\pi x) dx$
34.  $\int_0^{\pi/2} \sin x dx$

 In Exercises 35–38, use a computer algebra system and the error formulas to find  $n$  such that the error in the approximation of the definite integral is less than 0.00001 using (a) the Trapezoidal Rule and (b) Simpson's Rule.

35.  $\int_0^2 \sqrt{1+x} dx$
36.  $\int_0^2 (x+1)^{2/3} dx$
37.  $\int_0^1 \tan x^2 dx$
38.  $\int_0^1 \sin x^2 dx$

39. Approximate the area of the shaded region using (a) the Trapezoidal Rule and (b) Simpson's Rule with  $n = 4$ .




Figure for 39





Figure for 40

40. Approximate the area of the shaded region using (a) the Trapezoidal Rule and (b) Simpson's Rule with  $n = 8$ .

 41. **Programming** Write a program for a graphing utility to approximate a definite integral using the Trapezoidal Rule and Simpson's Rule. Start with the program written in Section 4.3, Exercises 59–62, and note that the Trapezoidal Rule can be written as  $T(n) = \frac{1}{2}[L(n) + R(n)]$  and Simpson's Rule can be written as

$$S(n) = \frac{1}{3}[T(n/2) + 2M(n/2)].$$

[Recall that  $L(n)$ ,  $M(n)$ , and  $R(n)$  represent the Riemann sums using the left-hand endpoints, midpoints, and right-hand endpoints of subintervals of equal width.]



**Programming** In Exercises 42–44, use the program in Exercise 41 to approximate the definite integral and complete the table.

$n$	$L(n)$	$M(n)$	$R(n)$	$T(n)$	$S(n)$
4					
8					
10					
12					
16					
20					

42.  $\int_0^4 \sqrt{2+3x^2} dx$     43.  $\int_0^1 \sqrt{1-x^2} dx$     44.  $\int_0^4 \sin \sqrt{x} dx$

45. **Area** Use Simpson's Rule with  $n = 14$  to approximate the area of the region bounded by the graphs of  $y = \sqrt{x} \cos x$ ,  $y = 0$ ,  $x = 0$ , and  $x = \pi/2$ .

46. **Circumference** The elliptic integral

$$8\sqrt{3} \int_0^{\pi/2} \sqrt{1 - \frac{2}{3} \sin^2 \theta} d\theta$$

gives the circumference of an ellipse. Use Simpson's Rule with  $n = 8$  to approximate the circumference.

47. **Work** To determine the size of the motor required to operate a press, a company must know the amount of work done when the press moves an object linearly 5 feet. The variable force to move the object is

$$F(x) = 100x \sqrt{125 - x^3}$$

where  $F$  is given in pounds and  $x$  gives the position of the unit in feet. Use Simpson's Rule with  $n = 12$  to approximate the work  $W$  (in foot-pounds) done through one cycle if

$$W = \int_0^5 F(x) dx.$$

48. The table lists several measurements gathered in an experiment to approximate an unknown continuous function  $y = f(x)$ .

(a) Approximate the integral  $\int_0^2 f(x) dx$  using the Trapezoidal Rule and Simpson's Rule.

$x$	0.00	0.25	0.50	0.75	1.00
$y$	4.32	4.36	4.58	5.79	6.14

$x$	1.25	1.50	1.75	2.00
$y$	7.25	7.64	8.08	8.14

(b) Use a graphing utility to find a model of the form  $y = ax^3 + bx^2 + cx + d$  for the data. Integrate the resulting polynomial over  $[0, 2]$  and compare the result with part (a).


**Approximation of  $\pi$**  In Exercises 49 and 50, use Simpson's Rule with  $n = 6$  to approximate  $\pi$  using the given equation. (In Section 5.7, you will be able to evaluate the integral using inverse trigonometric functions.)

49.  $\pi = \int_0^{1/2} \frac{6}{\sqrt{1-x^2}} dx$     50.  $\pi = \int_0^1 \frac{4}{1+x^2} dx$

**Area** In Exercises 51 and 52, use the Trapezoidal Rule to estimate the number of square meters of land in a lot where  $x$  and  $y$  are measured in meters, as shown in the figures. The land is bounded by a stream and two straight roads that meet at right angles.


51.

$x$	$y$
0	125
100	125
200	120
300	112
400	90
500	90
600	95
700	88
800	75
900	35
1000	0



52.

$x$	$y$
0	75
10	81
20	84
30	76
40	67
50	68
60	69
70	72
80	68
90	56
100	42
110	23
120	0



53. Prove that Simpson's Rule is exact when approximating the integral of a cubic polynomial function, and demonstrate the result for  $\int_0^1 x^3 dx$ ,  $n = 2$ .

54. Use Simpson's Rule with  $n = 10$  and a computer algebra system to approximate  $t$  in the integral equation


$$\int_0^t \sin \sqrt{x} dx = 2.$$

55. Prove that you can find a polynomial  $p(x) = Ax^2 + Bx + C$  that passes through any three points  $(x_1, y_1)$ ,  $(x_2, y_2)$ , and  $(x_3, y_3)$ , where the  $x_i$ 's are distinct.

## Review Exercises for Chapter 4

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.

In Exercises 1 and 2, use the graph of  $f'$  to sketch a graph of  $f$ . To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).



In Exercises 3–8, find the indefinite integral.

3.  $\int (2x^2 + x - 1) dx$

4.  $\int \frac{2}{\sqrt[3]{3x}} dx$

5.  $\int \frac{x^3 + 1}{x^2} dx$


6.  $\int \frac{x^3 - 2x^2 + 1}{x^2} dx$

7.  $\int (4x - 3 \sin x) dx$

8.  $\int (5 \cos x - 2 \sec^2 x) dx$



9. Find the particular solution of the differential equation  $f'(x) = -2x$  whose graph passes through the point  $(-1, 1)$ .

10. Find the particular solution of the differential equation  $f''(x) = 6(x - 1)$  whose graph passes through the point  $(2, 1)$  and is tangent to the line  $3x - y - 5 = 0$  at that point.

 **Slope Fields** In Exercises 11 and 12, a differential equation, a point, and a slope field are given. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the given point. (To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).) (b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution.

11.  $\frac{dy}{dx} = 2x - 4$ ,  $(4, -2)$

12.  $\frac{dy}{dx} = \frac{1}{2}x^2 - 2x$ ,  $(6, 2)$



13. **Velocity and Acceleration** An airplane taking off from a runway travels 3600 feet before lifting off. The airplane starts from rest, moves with constant acceleration, and makes the run in 30 seconds. With what speed does it lift off?

14. **Velocity and Acceleration** The speed of a car traveling in a straight line is reduced from 45 to 30 miles per hour in a distance of 264 feet. Find the distance in which the car can be brought to rest from 30 miles per hour, assuming the same constant deceleration.

15. **Velocity and Acceleration** A ball is thrown vertically upward from ground level with an initial velocity of 96 feet per second.

(a) How long will it take the ball to rise to its maximum height?

(b) What is the maximum height?

(c) When is the velocity of the ball one-half the initial velocity?

(d) What is the height of the ball when its velocity is one-half the initial velocity?

16. **Velocity and Acceleration** Repeat Exercise 15 for an initial velocity of 40 meters per second.

In Exercises 17–20, use sigma notation to write the sum.

17.  $\frac{1}{4(1)} + \frac{1}{4(2)} + \frac{1}{4(3)} + \cdots + \frac{1}{4(8)}$

18.  $\frac{1+2}{2(1)} + \frac{2+2}{2(2)} + \frac{3+2}{2(3)} + \cdots + \frac{12+2}{2(12)}$

19.  $\left(\frac{3}{n}\right)\left(\frac{1+1}{n}\right)^2 + \left(\frac{3}{n}\right)\left(\frac{2+1}{n}\right)^2 + \cdots + \left(\frac{3}{n}\right)\left(\frac{n+1}{n}\right)^2$

20.  $3\left(2 + \frac{4}{n}\right) + \cdots + 3n\left(2 + \frac{(n+1)^2}{n}\right)$

In Exercises 21–24, use the properties of summation and Theorem 4.2 to evaluate the sum.

21.  $\sum_{i=1}^{10} 3i$

22.  $\sum_{i=1}^{20} (4i - 1)$

23.  $\sum_{i=1}^{20} (i + 1)^2$

24.  $\sum_{i=1}^{12} i(i^2 - 1)$

25. Write in sigma notation (a) the sum of the first ten positive odd integers, (b) the sum of the cubes of the first  $n$  positive integers, and (c)  $6 + 10 + 14 + 18 + \cdots + 42$ .

26. Evaluate each sum for  $x_1 = 2$ ,  $x_2 = -1$ ,  $x_3 = 5$ ,  $x_4 = 3$ , and  $x_5 = 7$ .

(a)  $\frac{1}{5} \sum_{i=1}^5 x_i$


(b)  $\sum_{i=1}^5 \frac{1}{x_i}$

(c)  $\sum_{i=1}^5 (2x_i - x_i^2)$


(d)  $\sum_{i=2}^5 (x_i - x_{i-1})$

In Exercises 27 and 28, use upper and lower sums to approximate the area of the region using the indicated number of subintervals of equal width.

27.  $y = \frac{10}{x^2 + 1}$



28.  $y = 9 - \frac{1}{4}x^2$



In Exercises 29–32, use the limit process to find the area of the region between the graph of the function and the  $x$ -axis over the given interval. Sketch the region.

29.  $y = 6 - x$ ,  $[0, 4]$

30.  $y = x^2 + 3$ ,  $[0, 2]$

31.  $y = 5 - x^2$ ,  $[-2, 1]$

32.  $y = \frac{1}{4}x^3$ ,  $[2, 4]$

33. Use the limit process to find the area of the region bounded by  $x = 5y - y^2$ ,  $x = 0$ ,  $y = 2$ , and  $y = 5$ .

34. Consider the region bounded by  $y = mx$ ,  $y = 0$ ,  $x = 0$ , and  $x = b$ .

- Find the upper and lower sums to approximate the area of the region when  $\Delta x = b/4$ .
- Find the upper and lower sums to approximate the area of the region when  $\Delta x = b/n$ .
- Find the area of the region by letting  $n$  approach infinity in both sums in part (b). Show that in each case you obtain the formula for the area of a triangle.

In Exercises 35 and 36, write the limit as a definite integral on the interval  $[a, b]$ , where  $c_i$  is any point in the  $i$ th subinterval.

Limit

Interval

35.  $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n (2c_i - 3) \Delta x_i$


$[4, 6]$

36.  $\lim_{\|\Delta\| \rightarrow 0} \sum_{i=1}^n 3c_i(9 - c_i^2) \Delta x_i$


$[1, 3]$

In Exercises 37 and 38, set up a definite integral that yields the area of the region. (Do not evaluate the integral.)

37.  $f(x) = 3x + 6$



38.  $f(x) = 9 - x^2$



In Exercises 39 and 40, sketch the region whose area is given by the definite integral. Then use a geometric formula to evaluate the integral.

39.  $\int_0^5 (5 - |x - 5|) dx$

40.  $\int_{-4}^4 \sqrt{16 - x^2} dx$

41. Given  $\int_2^6 f(x) dx = 10$  and  $\int_2^6 g(x) dx = 3$ , evaluate

(a)  $\int_2^6 [f(x) + g(x)] dx$ .

(b)  $\int_2^6 [f(x) - g(x)] dx$ .

(c)  $\int_2^6 [2f(x) - 3g(x)] dx$ .

(d)  $\int_2^6 5f(x) dx$ .

42. Given  $\int_0^3 f(x) dx = 4$  and  $\int_3^6 f(x) dx = -1$ , evaluate

(a)  $\int_0^6 f(x) dx$ .

(b)  $\int_6^3 f(x) dx$ .

(c)  $\int_4^4 f(x) dx$ .

(d)  $\int_3^6 -10f(x) dx$ .

In Exercises 43–50, use the Fundamental Theorem of Calculus to evaluate the definite integral.

43.  $\int_0^4 (2 + x) dx$

44.  $\int_{-1}^1 (t^2 + 2) dt$

45.  $\int_{-1}^1 (4t^3 - 2t) dt$

46.  $\int_{-2}^2 (x^4 + 2x^2 - 5) dx$

47.  $\int_4^9 x\sqrt{x} dx$

48.  $\int_1^2 \left( \frac{1}{x^2} - \frac{1}{x^3} \right) dx$

49.  $\int_0^{3\pi/4} \sin \theta d\theta$

50.  $\int_{-\pi/4}^{\pi/4} \sec^2 t dt$

In Exercises 51–56, sketch the graph of the region whose area is given by the integral, and find the area.

51.  $\int_1^3 (2x - 1) dx$

52.  $\int_0^2 (x + 4) dx$

53.  $\int_3^4 (x^2 - 9) dx$


54.  $\int_{-1}^2 (-x^2 + x + 2) dx$

55.  $\int_0^1 (x - x^3) dx$


56.  $\int_0^1 \sqrt{x}(1 - x) dx$

In Exercises 57 and 58, determine the area of the given region.

57.  $y = \sin x$



58.  $y = x + \cos x$



In Exercises 59 and 60, sketch the region bounded by the graphs of the equations, and determine its area.

59.  $y = \frac{4}{\sqrt{x}}$ ,  $y = 0$ ,  $x = 1$ ,  $x = 9$

60.  $y = \sec^2 x$ ,  $y = 0$ ,  $x = 0$ ,  $x = \frac{\pi}{3}$

In Exercises 61 and 62, find the average value of the function over the given interval. Find the values of  $x$  at which the function assumes its average value, and graph the function.

61.  $f(x) = \frac{1}{\sqrt{x}}$ ,  $[4, 9]$

62.  $f(x) = x^3$ ,  $[0, 2]$

In Exercises 63–66, use the Second Fundamental Theorem of Calculus to find  $F'(x)$ .

63.  $F(x) = \int_0^x t^2 \sqrt{1+t^3} dt$

64.  $F(x) = \int_1^x \frac{1}{t^2} dt$

65.  $F(x) = \int_{-3}^x (t^2 + 3t + 2) dt$

66.  $F(x) = \int_0^x \csc^2 t dt$

In Exercises 67–80, find the indefinite integral.

67.  $\int (x^2 + 1)^3 dx$

68.  $\int \left(x + \frac{1}{x}\right)^2 dx$

69.  $\int \frac{x^2}{\sqrt{x^3 + 3}} dx$

70.  $\int x^2 \sqrt{x^3 + 3} dx$

71.  $\int x(1 - 3x^2)^4 dx$

72.  $\int \frac{x+3}{(x^2 + 6x - 5)^2} dx$

73.  $\int \sin^3 x \cos x dx$

74.  $\int x \sin 3x^2 dx$

75.  $\int \frac{\sin \theta}{\sqrt{1 - \cos \theta}} d\theta$

76.  $\int \frac{\cos x}{\sqrt{\sin x}} dx$

77.  $\int \tan^n x \sec^2 x dx$ ,  $n \neq -1$

78.  $\int \sec 2x \tan 2x dx$

79.  $\int (1 + \sec \pi x)^2 \sec \pi x \tan \pi x dx$

80.  $\int \cot^4 \alpha \csc^2 \alpha d\alpha$

In Exercises 81–88, evaluate the definite integral. Use a graphing utility to verify your result.

81.  $\int_{-1}^2 x(x^2 - 4) dx$

82.  $\int_0^1 x^2(x^3 + 1)^3 dx$

83.  $\int_0^3 \frac{1}{\sqrt{1+x}} dx$

84.  $\int_3^6 \frac{x}{3\sqrt{x^2 - 8}} dx$

85.  $2\pi \int_0^1 (y+1)\sqrt{1-y} dy$

86.  $2\pi \int_{-1}^0 x^2 \sqrt{x+1} dx$



87.  $\int_0^\pi \cos \frac{x}{2} dx$

88.  $\int_{-\pi/4}^{\pi/4} \sin 2x dx$

**Slope Fields** In Exercises 89 and 90, a differential equation, a point, and a slope field are given. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the given point. (To print an enlarged copy of the graph, go to the website [www.mathgraphs.com](http://www.mathgraphs.com).)

(b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution.



89.  $\frac{dy}{dx} = x\sqrt{9-x^2}$ ,  $(0, -4)$     90.  $\frac{dy}{dx} = -\frac{1}{2}x \sin(x^2)$ ,  $(0, 0)$



In Exercises 91 and 92, find the area of the region. Use a graphing utility to verify your result.

91.  $\int_1^9 x \sqrt[3]{x-1} dx$

92.  $\int_0^{\pi/2} [\cos x + \sin(2x)] dx$



93. **Fuel Cost** Gasoline is increasing in price according to the equation  $p = 1.20 + 0.04t$ , where  $p$  is the dollar price per gallon and  $t$  is the time in years, with  $t = 0$  representing 1990. An automobile is driven 15,000 miles a year and gets  $M$  miles per gallon. The annual fuel cost is

$$C = \frac{15,000}{M} \int_t^{t+1} p dt.$$

Estimate the annual fuel cost in (a) 2000 and (b) 2005.

94. **Respiratory Cycle** After exercising for a few minutes, a person has a respiratory cycle for which the rate of air intake is

$$v = 1.75 \sin \frac{\pi t}{2}.$$

Find the volume, in liters, of air inhaled during one cycle by integrating the function over the interval  $[0, 2]$ .



In Exercises 95–98, use the Trapezoidal Rule and Simpson's Rule with  $n = 4$ , and use the integration capabilities of a graphing utility, to approximate the definite integral. Compare the results.

95.  $\int_1^2 \frac{1}{1+x^3} dx$

96.  $\int_0^1 \frac{x^{3/2}}{3-x^2} dx$

97.  $\int_0^{\pi/2} \sqrt{x} \cos x dx$

98.  $\int_0^\pi \sqrt{1+\sin^2 x} dx$

## P.S.


## Problem Solving

See [www.CalcChat.com](http://www.CalcChat.com) for worked-out solutions to odd-numbered exercises.


1. Let  $L(x) = \int_1^x \frac{1}{t} dt$ ,  $x > 0$ .

(a) Find  $L(1)$ .

(b) Find  $L'(x)$  and  $L'(1)$ .

 (c) Use a graphing utility to approximate the value of  $x$  (to three decimal places) for which  $L(x) = 1$ .

(d) Prove that  $L(x_1 x_2) = L(x_1) + L(x_2)$  for all positive values of  $x_1$  and  $x_2$ .

 2. Let  $F(x) = \int_2^x \sin t^2 dt$ .


(a) Use a graphing utility to complete the table.

$x$	0	1.0	1.5	1.9	2.0
$F(x)$					
$x$	2.1	2.5	3.0	4.0	5.0
$F(x)$					

(b) Let  $G(x) = \frac{1}{x-2} F(x) = \frac{1}{x-2} \int_2^x \sin t^2 dt$ . Use a graphing utility to complete the table and estimate  $\lim_{x \rightarrow 2} G(x)$ .

$x$	1.9	1.95	1.99	2.01	2.1
$G(x)$					

(c) Use the definition of the derivative to find the exact value of the limit  $\lim_{x \rightarrow 2} G(x)$ .

 In Exercises 3 and 4, (a) write the area under the graph of the given function defined on the given interval as a limit. Then use a computer algebra system to (b) evaluate the sum in part (a), and (c) evaluate the limit using the result of part (b).

3.  $y = x^4 - 4x^3 + 4x^2$ ,  $[0, 2]$

(Hint:  $\sum_{i=1}^n i^4 = \frac{n(n+1)(2n+1)(3n^2+3n-1)}{30}$ )

4.  $y = \frac{1}{2}x^5 + 2x^3$ ,  $[0, 2]$

(Hint:  $\sum_{i=1}^n i^5 = \frac{n^2(n+1)^2(2n^2+2n-1)}{12}$ )

5. The **Fresnel function**  $S$  is defined by the integral

$$S(x) = \int_0^x \sin\left(\frac{\pi t^2}{2}\right) dt.$$

(a) Graph the function  $y = \sin\left(\frac{\pi x^2}{2}\right)$  on the interval  $[0, 3]$ .

(b) Use the graph in part (a) to sketch the graph of  $S$  on the interval  $[0, 3]$ .

(c) Locate all relative extrema of  $S$  on the interval  $(0, 3)$ .

(d) Locate all points of inflection of  $S$  on the interval  $(0, 3)$ .

6. The **Two-Point Gaussian Quadrature Approximation** for  $f$  is


$$\int_{-1}^1 f(x) dx \approx f\left(-\frac{1}{\sqrt{3}}\right) + f\left(\frac{1}{\sqrt{3}}\right).$$

(a) Use this formula to approximate  $\int_{-1}^1 \cos x dx$ . Find the error of the approximation.

(b) Use this formula to approximate  $\int_{-1}^1 \frac{1}{1+x^2} dx$ .

(c) Prove that the Two-Point Gaussian Quadrature Approximation is exact for all polynomials of degree 3 or less.

7. Archimedes showed that the area of a parabolic arch is equal to  $\frac{2}{3}$  the product of the base and the height (see figure).



(a) Graph the parabolic arch bounded by  $y = 9 - x^2$  and the  $x$ -axis. Use an appropriate integral to find the area  $A$ .

(b) Find the base and height of the arch and verify Archimedes' formula.

(c) Prove Archimedes' formula for a general parabola.

8. Galileo Galilei (1564–1642) stated the following proposition concerning falling objects:

*The time in which any space is traversed by a uniformly accelerating body is equal to the time in which that same space would be traversed by the same body moving at a uniform speed whose value is the mean of the highest speed of the accelerating body and the speed just before acceleration began.*

Use the techniques of this chapter to verify this proposition.

9. The graph of the function  $f$  consists of the three line segments joining the points  $(0, 0)$ ,  $(2, -2)$ ,  $(6, 2)$ , and  $(8, 3)$ . The function  $F$  is defined by the integral

$$F(x) = \int_0^x f(t) dt.$$

(a) Sketch the graph of  $f$ .

(b) Complete the table.

$x$	0	1	2	3	4	5	6	7	8
$F(x)$									

(c) Find the extrema of  $F$  on the interval  $[0, 8]$ .

(d) Determine all points of inflection of  $F$  on the interval  $(0, 8)$ .



10. A car is traveling in a straight line for 1 hour. Its velocity  $v$  in miles per hour at six-minute intervals is shown in the table.

$t$ (hours)	0	0.1	0.2	0.3	0.4	0.5
$v$ (mi/h)	0	10	20	40	60	50

$t$ (hours)	0.6	0.7	0.8	0.9	1.0
$v$ (mi/h)	40	35	40	50	65

- (a) Produce a reasonable graph of the velocity function  $v$  by graphing these points and connecting them with a smooth curve.
- (b) Find the open intervals over which the acceleration  $a$  is positive.
- (c) Find the average acceleration of the car (in miles per hour squared) over the interval  $[0, 0.4]$ .
- (d) What does the integral  $\int_0^1 v(t) dt$  signify? Approximate this integral using the Trapezoidal Rule with five subintervals.
- (e) Approximate the acceleration at  $t = 0.8$ .

11. Prove  $\int_0^x f(t)(x-t) dt = \int_0^x \left( \int_0^t f(v) dv \right) dt$ .

12. Prove  $\int_a^b f(x)f'(x) dx = \frac{1}{2}([f(b)]^2 - [f(a)]^2)$ .

13. Use an appropriate Riemann sum to evaluate the limit

$$\lim_{n \rightarrow \infty} \frac{\sqrt{1} + \sqrt{2} + \sqrt{3} + \cdots + \sqrt{n}}{n^{3/2}}.$$

14. Use an appropriate Riemann sum to evaluate the limit

$$\lim_{n \rightarrow \infty} \frac{1^5 + 2^5 + 3^5 + \cdots + n^5}{n^6}.$$

15. Suppose that  $f$  is integrable on  $[a, b]$  and  $0 < m \leq f(x) \leq M$  for all  $x$  in the interval  $[a, b]$ . Prove that

$$m(a-b) \leq \int_a^b f(x) dx \leq M(b-a).$$

Use this result to estimate  $\int_0^1 \sqrt{1+x^4} dx$ .

16. Let  $f$  be continuous on the interval  $[0, b]$  where  $f(x) + f(b-x) \neq 0$  on  $[0, b]$ .

(a) Show that  $\int_0^b \frac{f(x)}{f(x) + f(b-x)} dx = \frac{b}{2}$ .

- (b) Use the result in part (a) to evaluate

$$\int_0^1 \frac{\sin x}{\sin(1-x) + \sin x} dx.$$

- (c) Use the result in part (a) to evaluate

$$\int_0^3 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{3-x}} dx.$$

17. Verify that

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

by showing the following.

(a)  $(1+i)^3 - i^3 = 3i^2 + 3i + 1$

(b)  $(n+1)^3 = \sum_{i=1}^n (3i^2 + 3i + 1) + 1$

(c)  $\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$

18. Prove that if  $f$  is a continuous function on a closed interval  $[a, b]$ , then


$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$

19. Let

$$I = \int_0^4 f(x) dx$$

where  $f$  is shown in the figure. Let  $L(n)$  and  $R(n)$  represent the Riemann sums using the left-hand endpoints and right-hand endpoints of  $n$  subintervals of equal width. (Assume  $n$  is even.) Let  $T(n)$  and  $S(n)$  be the corresponding values of the Trapezoidal Rule and Simpson's Rule.

- (a) For any  $n$ , list  $L(n)$ ,  $R(n)$ ,  $T(n)$ , and  $I$  in increasing order.
- (b) Approximate  $S(4)$ .



20. The sine integral function

$$\text{Si}(x) = \int_0^x \frac{\sin t}{t} dt$$

is often used in engineering. The function  $f(t) = \frac{\sin t}{t}$  is not defined at  $t = 0$ , but its limit is 1 as  $t \rightarrow 0$ . So, define  $f(0) = 1$ .

Then  $f$  is continuous everywhere.



- (a) Use a graphing utility to graph  $\text{Si}(x)$ .
- (b) At what values of  $x$  does  $\text{Si}(x)$  have relative maxima?
- (c) Find the coordinates of the first inflection point where  $x > 0$ .
- (d) Decide whether  $\text{Si}(x)$  has any horizontal asymptotes. If so, identify each.